

A CHARACTERIZATION OF SOLUTIONS OF AFFINE VARIATIONAL INEQUALITY DEFINED BY SECOND-ORDER CONE

SANGHO KUM*

ABSTRACT. A variational inequality defined by the second-order cone is considered, and it is shown that a necessary and sufficient condition for a solution of the variational inequality holds under some regularity condition.

1. Introduction and preliminaries

Recently, many authors have studied optimization problems over second-order cones and their applications to nonlinear optimization problems [1, 2, 6]. Such investigation has a strong motivation since many nonlinear optimization problems can be relaxed to optimization problems over second-order cones. It is well-known that variational inequality problems are closely related to optimization problems. Very recently, Auslender [3] studied algorithms for variational inequalities over the cone of semidefinite positive symmetric matrices and over the second-order cone.

The purpose of this paper is to give a complete characterization of solutions for a variational inequality problem defined by the second-order cone. We prove that a necessary and sufficient condition for a solution of the variational inequality holds under some regularity condition. We recall in this section some notations and basic results which will be used in next section.

Received August 6, 2007.

2000 Mathematics Subject Classification: 47J20, 65K10.

Key words and phrases: variational inequality, second-order cone, regularity condition.

This work was supported by the research grant of the Chungbuk National University in 2006.

For a subset $D \subset \mathbb{R}^n$, the closure of D will be denoted by $\text{cl}D$ and the convex hull of D by $\text{co}D$. Let $h : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$ be a proper, lower semi-continuous convex function. The conjugate function of h , $h^* : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$, is defined by

$$f^*(v) := \sup\{v(x) - h(x) \mid x \in \text{dom } h\},$$

where $\text{dom } h := \{x \in \mathbb{R}^n \mid h(x) < +\infty\}$ is the effective domain of h . The epigraph of h is defined by

$$\text{epi } h := \{(x, r) \in \mathbb{R}^n \times \mathbb{R} \mid x \in \text{dom } h, h(x) \leq r\}.$$

The set (possibly empty)

$$\partial h(a) := \{v \in \mathbb{R}^n \mid h(x) - h(a) \geq v(x - a), \forall x \in \text{dom } h\}$$

is the subdifferential of the convex function h at $a \in \text{dom } h$. For a closed convex subset D of \mathbb{R}^n , the indicator function δ_D is defined as $\delta_D(x) = 0$ if $x \in D$ and $\delta_D(x) = +\infty$ if $x \notin D$. Then $\partial \delta_D(x) = N_D(x)$, which is known as the normal cone of D of x .

For a proper lower semicontinuous convex functions $g : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{+\infty\}$, the lower semicontinuous convex hull of g is denoted by cl cog . That is, $\text{epi}(\text{cl cog}) = \text{cl co}(\text{epi } g)$. For details, see [7]. Let g_i , $i \in I$ (where I is an arbitrary index set) be proper lower semicontinuous convex functions. It is well known (see [7]) that if $\sup_{i \in I} g_i$ is proper, then

$$\left(\sup_{i \in I} g_i \right)^* = \text{cl co} \left(\inf_{i \in I} g_i^* \right).$$

Thus we can check that

$$(1.1) \quad \text{epi} \left(\sup_{i \in I} g_i \right)^* = \text{cl co} \left(\bigcup_{i \in I} \text{epi } g_i^* \right).$$

2. A characterization of solutions

In this section, we introduce a variational inequality defined by the second-order cone, and then we prove that a necessary and sufficient condition for a point to be a solution of the inequality holds under a regularity condition. The condition gives a complete characterization of solutions of $\text{Aff}(\text{VI})$.

Let $Q := \{x \in \mathbb{R}^n \mid \|Hx + b\| \leq c^t x + d\}$, where H is an $(m - 1) \times n$ matrix, $b \in \mathbb{R}^{m-1}$, $c \in \mathbb{R}^n$, $d \in \mathbb{R}$ and $\|z\| = \sqrt{z^t z}$, $z \in \mathbb{R}^{m-1}$. Suppose that $Q \neq \emptyset$. Let $K = \{(y, t)^t \in \mathbb{R}^{m-1} \times \mathbb{R} \mid \|y\| \leq t\}$, that

is, K is a second-order cone in \mathbb{R}^m . Then K is self-dual, that is, $K = \{z \in \mathbb{R}^m \mid k^t z \geq 0 \ \forall z \in K\}$. Consider the following affine variational inequality defined by the second-order cone K .

Aff(VI) : Find $\bar{x} \in Q$ such that $\langle M\bar{x}, x - \bar{x} \rangle \geq 0 \quad \forall x \in Q$,

where $M \in \mathbb{R}^{n \times n}$ is a given matrix. Note that

$$(2.1) \quad x \in Q \Leftrightarrow \begin{pmatrix} Hx + b \\ c^t x + d \end{pmatrix} \in K.$$

Thus Aff(VI) is a variational inequality defined by the second-order cone K . We begin with the following lemma.

LEMMA 2.1. *Let $\bar{x} \in Q$. Then we have the following:*

$$u \in N_Q(\bar{x}) \iff (u, u^t \bar{x}) \in \text{cl} \left[\bigcup_{\lambda \in K} \left\{ \begin{pmatrix} -\left(\frac{H}{c^t}\right)^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda \end{pmatrix} \right\} + \{0\} \times \mathbb{R}^+ \right].$$

Proof. Let $g(x) = -\begin{pmatrix} Hx + b \\ c^t x + d \end{pmatrix}$. For any $\lambda \in K$, we get

$$\begin{aligned} (\lambda^t g)^*(v) &= \sup \{ v^t x - \lambda^t g(x) \mid x \in \mathbb{R}^n \} \\ &= \sup \left\{ \left[v + \begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda \right]^t x \mid x \in \mathbb{R}^n \right\} + \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda \\ &= \begin{cases} \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda & \text{if } v = -\begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda \\ +\infty & \text{if } v \neq -\begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda, \end{cases} \end{aligned}$$

and hence

$$\begin{aligned} \text{epi}(\lambda^t g)^* &= \left\{ \begin{pmatrix} -\left(\frac{H}{c^t}\right)^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda + r \end{pmatrix} \mid r \geq 0 \right\} \\ &= \begin{pmatrix} -\left(\frac{H}{c^t}\right)^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda \end{pmatrix} + \{0\} \times \mathbb{R}^+. \end{aligned}$$

Thus we have

$$\bigcup_{\lambda \in K} \text{epi}(\lambda^t g)^* = \bigcup_{\lambda \in K} \left\{ \begin{pmatrix} -\left(\frac{H}{c^t}\right)^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda \end{pmatrix} \right\} + \{0\} \times \mathbb{R}^+.$$

Since $\bar{x} \in Q$, we see

$$\begin{aligned} u \in N_Q(\bar{x}) &\iff u^t(x - \bar{x}) \leq 0 \quad \forall x \in Q \\ &\iff \sup_{x \in Q} u^t x = u^t \bar{x} \\ &\iff \delta_Q^*(u) = u^t \bar{x} \\ &\iff (u, u^t \bar{x}) \in \text{epi } \delta_Q^*. \end{aligned}$$

As $\delta_Q(x) = \sup_{\lambda \in K} \lambda^t g(x)$ and $\bigcup_{\lambda \in K} \text{epi}(\lambda^t g)^*$ is convex [4], it follows from (1.1) that

$$\text{epi } \delta_Q^* = \text{cl co} \left[\bigcup_{\lambda \in K} \text{epi}(\lambda^t g)^* \right] = \text{cl} \left[\bigcup_{\lambda \in K} \text{epi}(\lambda^t g)^* \right].$$

This completes the proof. \square

Now we present the main result of this note.

THEOREM 2.2. *Assume that $\bigcup_{\lambda \in K} \left\{ \left(- \begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda \right) \right\} + \{0\} \times \mathbb{R}^+$ is closed. Then $\bar{x} \in \mathbb{R}^n$ is a solution of Aff(VI) if and only if $\begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} \in K$, and there exists $\lambda \in K$ such that*

$$M\bar{x} = \begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda \quad \text{and} \quad \lambda^t \begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} = 0.$$

Proof. Let us denote by $\text{sol}(\text{Aff(VI)})$ the solution set of Aff(VI) . Then we have the following equivalences:

$$\bar{x} \in \text{sol}(\text{Aff(VI)}) \iff \bar{x} \in Q \quad \text{and} \quad -M\bar{x} \in N_Q(\bar{x})$$

$$\begin{aligned}
&\iff \bar{x} \in Q \text{ and } \left[\exists \lambda \in K \text{ and } \exists \alpha \geq 0 \text{ satisfying} \right. \\
&\quad \left. (-M\bar{x}, -\bar{x}^t M^t \bar{x}) = \left(-\begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda, \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda + \alpha \right) \right. \\
&\quad \left. \text{i.e., } -M\bar{x} = -\begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda, -\bar{x}^t M^t \bar{x} = \begin{pmatrix} b \\ d \end{pmatrix}^t \lambda + \alpha \right] \\
&\iff \begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} \in K \text{ and } \left[\exists \lambda \in K \text{ and } \exists \alpha \geq 0 \text{ such that} \right. \\
&\quad \left. M\bar{x} = \begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda \text{ and } -\lambda^t \begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} = \alpha \right] \\
&\iff \begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} \in K \text{ and } \left[\exists \lambda \in K \text{ such that} \right. \\
&\quad \left. M\bar{x} = \begin{pmatrix} H \\ c^t \end{pmatrix}^t \lambda \text{ and } \lambda^t \begin{pmatrix} H\bar{x} + b \\ c^t \bar{x} + d \end{pmatrix} = 0. \right]
\end{aligned}$$

The second and third equivalences come from Lemma 2.1 and (2.1), respectively. This completes the proof. \square

References

- [1] F. Alizadeh and D. Goldfarb, *Second-order cone programming*, Math. Prog. **95** (2003), 3-51.
- [2] E. D. Andersen, C. Roos and T. Terlaky, *Notes on duality in second order and p -order cone optimization*, Optimization **51** (2002), 627-643.
- [3] A. Auslender, *Variational inequalities over the cone of semidefinite positive symmetric matrices and over the Lorentz cone*, Optim. Meth. Software **18** (2003), 359-376.
- [4] V. Jeyakumar, A. M. Rubinov, B. M. Glover and Y. Ishizuka, *Inequality systems and global optimization*, J. Math. Anal. Appl. **202** (1996), 900-919.
- [5] V. Jeyakumar, *Constraint qualifications characterizing Lagrangian Duality in convex optimization*, submitted.
- [6] M. S. Lobo, L. Vandenberghe, S. Boyd and H. Lebret, *Applications of second-order cone programming*, Linear Algebra Appl. **284** (1998), 193-228.
- [7] C. Zalinescu, *Convex Analysis in General Vector Space*, World Scientific Publishing Co. Pte. Ltd, Singapore, 2002.

*

Department of Mathematics Education
Chungbuk National University
Cheongju 361-763, Republic of Korea
E-mail: shkum@chungbuk.ac.kr