

FUNCTIONAL EQUATIONS ASSOCIATED WITH INNER PRODUCT SPACES

CHOONKIL PARK*, JAE SUNG HUH**, WON JUNE MIN***, DONG
HOON NAM****, AND SEUNG HYEON ROH*****

ABSTRACT. In [7], Th.M. Rassias proved that the norm defined over a real vector space V is induced by an inner product if and only if for a fixed integer $n \geq 2$

$$n \left\| \frac{1}{n} \sum_{i=1}^n x_i \right\|^2 + \sum_{i=1}^n \left\| x_i - \frac{1}{n} \sum_{j=1}^n x_j \right\|^2 = \sum_{i=1}^n \|x_i\|^2$$

holds for all $x_1, \dots, x_n \in V$.

Let V, W be real vector spaces. It is shown that if a mapping $f : V \rightarrow W$ satisfies

$$(0.1) \quad n f \left(\frac{1}{n} \sum_{i=1}^n x_i \right) + \sum_{i=1}^n f \left(x_i - \frac{1}{n} \sum_{j=1}^n x_j \right) \\ = \sum_{i=1}^n f(x_i)$$

for all $x_1, \dots, x_n \in V$, then the mapping $f : V \rightarrow W$ satisfies

$$(0.2) \quad 2f \left(\frac{x+y}{2} \right) + f \left(\frac{x-y}{2} \right) + f \left(\frac{y-x}{2} \right) \\ = f(x) + f(y)$$

for all $x, y \in V$.

Furthermore, we prove the generalized Hyers-Ulam stability of the functional equation (0.2) in real Banach spaces.

Received August 18, 2008; Accepted November 20, 2008.

2000 Mathematics Subject Classification: Primary 39B72, 46C05.

Key words and phrases: Jensen additive mapping, Jensen quadratic mapping, functional equation associated with inner product space, generalized Hyers-Ulam stability.

This work was supported by the R & E program of KOSEF in 2008.

1. Introduction

The stability problem of functional equations originated from a question of Ulam [15] concerning the stability of group homomorphisms. Hyers [5] gave a first affirmative partial answer to the question of Ulam for Banach spaces. Hyers' Theorem was generalized by Aoki [1] for additive mappings and by Th.M. Rassias [6] for linear mappings by considering an unbounded Cauchy difference. The paper of Th.M. Rassias [6] has provided a lot of influence in the development of what we call *generalized Hyers-Ulam stability* of functional equations. A generalization of the Th.M. Rassias theorem was obtained by Găvruta [4] by replacing the unbounded Cauchy difference by a general control function in the spirit of Th.M. Rassias' approach.

The functional equation

$$f(x+y) + f(x-y) = 2f(x) + 2f(y)$$

is called a *quadratic functional equation*. In particular, every solution of the quadratic functional equation is said to be a *quadratic mapping*. A generalized Hyers-Ulam stability problem for the quadratic functional equation was proved by Skof [14] for mappings $f : X \rightarrow Y$, where X is a normed space and Y is a Banach space. Cholewa [2] noticed that the theorem of Skof is still true if the relevant domain X is replaced by an Abelian group. In [3], Czerwik proved the generalized Hyers-Ulam stability of the quadratic functional equation. Several functional equations have been investigated in [8]–[13].

Throughout this paper, assume that n is a fixed integer greater than 1. Let X be a real normed vector space with norm $\|\cdot\|$, and Y a real Banach space with norm $\|\cdot\|$.

In this paper, we investigate the functional equation (0.2), and prove the generalized Hyers-Ulam stability of the functional equation (0.2) in real Banach spaces.

2. Jensen quadratic mappings associated with inner product spaces

We investigate the functional equations (0.1) and (0.2).

LEMMA 2.1. *Let V and W be real vector spaces. If a mapping $f : V \rightarrow W$ satisfies*

$$(2.1) \quad nf\left(\frac{1}{n}\sum_{i=1}^n x_i\right) + \sum_{i=1}^n f\left(x_i - \frac{1}{n}\sum_{j=1}^n x_j\right) = \sum_{i=1}^n f(x_i)$$

for all $x_1, \dots, x_n \in V$, then the mapping $f : V \rightarrow W$ satisfies

$$(2.2) \quad 2f\left(\frac{x+y}{2}\right) + f\left(\frac{x-y}{2}\right) + f\left(\frac{y-x}{2}\right) = f(x) + f(y)$$

for all $x, y \in V$.

Proof. Assume that $f : V \rightarrow W$ satisfies (2.1).

Letting $x_n = \frac{\sum_{i=1}^{n-1} x_i}{n-1}$, we get

$$(n-1)f\left(\frac{1}{n-1}\sum_{i=1}^{n-1} x_i\right) + \sum_{i=1}^{n-1} f\left(x_i - \frac{1}{n-1}\sum_{j=1}^{n-1} x_j\right) = \sum_{i=1}^{n-1} f(x_i)$$

for all $x_1, \dots, x_{n-1} \in V$. Applying continuously this method $n-3$ times, we get

$$2f\left(\frac{x_1+x_2}{2}\right) + f\left(\frac{x_1-x_2}{2}\right) + f\left(\frac{x_2-x_1}{2}\right) = f(x_1) + f(x_2)$$

for all $x_1, x_2 \in V$, as desired. $\square$

One can easily show that an even mapping $f : V \rightarrow W$ satisfies (2.2) if and only if the even mapping $f : V \rightarrow W$ is a Jensen quadratic mapping, i.e.,

$$2f\left(\frac{x+y}{2}\right) + 2f\left(\frac{x-y}{2}\right) = f(x) + f(y),$$

and that an odd mapping $f : V \rightarrow W$ satisfies (2.2) if and only if the odd mapping mapping $f : V \rightarrow W$ is a Jensen additive mapping, i.e.,

$$2f\left(\frac{x+y}{2}\right) = f(x) + f(y).$$

For a given mapping $f : X \rightarrow Y$, we define

$$Df(x, y) := 2f\left(\frac{x+y}{2}\right) + f\left(\frac{x-y}{2}\right) + f\left(\frac{y-x}{2}\right) - f(x) - f(y)$$

for all $x, y \in X$.

We prove the generalized Hyers-Ulam stability of the functional equation $Df(x, y) = 0$ in real Banach spaces: an even case.

THEOREM 2.2. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ such that*

$$(2.3) \quad \tilde{\varphi}(x, y) := \sum_{j=0}^{\infty} 4^j \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}\right) < \infty,$$

$$(2.4) \quad \|Df(x, y)\| \leq \varphi(x, y)$$

for all $x, y \in X$. Then there exists a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that

$$(2.5) \quad \|f(x) + f(-x) - Q(x)\| \leq \tilde{\varphi}(x, 0) + \tilde{\varphi}(-x, 0)$$

for all $x \in X$.

Proof. Letting $y = 0$ in (2.4), we get

$$(2.6) \quad \left\| 3f\left(\frac{x}{2}\right) + f\left(\frac{-x}{2}\right) - f(x) \right\| \leq \varphi(x, 0)$$

for all $x \in X$. Replacing x by $-x$ in (2.6), we get

$$(2.7) \quad \left\| 3f\left(\frac{-x}{2}\right) + f\left(\frac{x}{2}\right) - f(-x) \right\| \leq \varphi(-x, 0)$$

for all $x \in X$. Let $g(x) := f(x) + f(-x)$ for all $x \in X$. It follows from (2.6) and (2.7) that

$$(2.8) \quad \left\| g(x) - 4g\left(\frac{x}{2}\right) \right\| \leq \varphi(x, 0) + \varphi(-x, 0)$$

for all $x \in X$. Hence

$$(2.9) \quad \left\| 4^l g\left(\frac{x}{2^l}\right) - 4^m g\left(\frac{x}{2^m}\right) \right\| \leq \sum_{j=l}^{m-1} 4^j \varphi\left(\frac{x}{2^j}, 0\right) + \sum_{j=l}^{m-1} 4^j \varphi\left(-\frac{x}{2^j}, 0\right)$$

for all nonnegative integers m and l with $m > l$ and all $x \in X$. It follows from (2.3) and (2.9) that the sequence $\{4^k g(\frac{x}{2^k})\}$ is Cauchy for all $x \in X$. Since Y is complete, the sequence $\{4^k g(\frac{x}{2^k})\}$ converges. So one can define the mapping $Q : X \rightarrow Y$ by

$$Q(x) := \lim_{k \rightarrow \infty} 4^k g\left(\frac{x}{2^k}\right)$$

for all $x \in X$.

By (2.3) and (2.4),

$$\begin{aligned} \|DQ(x, y)\| &= \lim_{k \rightarrow \infty} 4^k \left\| Dg\left(\frac{x}{2^k}, \frac{y}{2^k}\right) \right\| \\ &\leq \lim_{k \rightarrow \infty} 4^k \left(\varphi\left(\frac{x}{2^k}, \frac{y}{2^k}\right) + \varphi\left(-\frac{x}{2^k}, -\frac{y}{2^k}\right) \right) = 0 \end{aligned}$$

for all $x, y \in X$. So $DQ(x, y) = 0$. Since $g : X \rightarrow Y$ is even, $Q : X \rightarrow Y$ is even. So the mapping $Q : X \rightarrow Y$ is Jensen quadratic. Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (2.9), we get (2.5). So there exists a Jensen quadratic mapping $Q : X \rightarrow Y$ satisfying (2.5).

Now, let $Q' : X \rightarrow Y$ be another Jensen quadratic mapping satisfying (2.5). Then we have

$$\begin{aligned} \|Q(x) - Q'(x)\| &= 4^q \left\| Q\left(\frac{x}{2^q}\right) - Q'\left(\frac{x}{2^q}\right) \right\| \\ &\leq 4^q \left\| Q\left(\frac{x}{2^q}\right) - f\left(\frac{x}{2^q}\right) - f\left(\frac{-x}{2^q}\right) \right\| \\ &\quad + 4^q \left\| Q'\left(\frac{x}{2^q}\right) - f\left(\frac{x}{2^q}\right) - f\left(\frac{-x}{2^q}\right) \right\| \\ &\leq 2 \cdot 4^q \tilde{\varphi}\left(\frac{x}{2^q}, 0\right) + 2 \cdot 4^q \tilde{\varphi}\left(\frac{-x}{2^q}, 0\right), \end{aligned}$$

which tends to zero as $q \rightarrow \infty$ for all $x \in X$. So we can conclude that $Q(x) = Q'(x)$ for all $x \in X$. This proves the uniqueness of Q . $\square$

COROLLARY 2.3. *Let $p > 2$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping such that*

$$(2.10) \quad \|Df(x, y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

for all $x, y \in X$. Then there exists a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that

$$\|f(x) + f(-x) - Q(x)\| \leq \frac{2^{p+1}\theta}{2^p - 4} \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 2.2 to get the desired result. $\square$

THEOREM 2.4. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ satisfying (2.4) such that*

$$(2.11) \quad \tilde{\varphi}(x, y) := \sum_{j=1}^{\infty} 4^{-j} \varphi(2^j x, 2^j y) < \infty$$

for all $x, y \in X$. Then there exists a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that

$$(2.12) \quad \|f(x) + f(-x) - Q(x)\| \leq \tilde{\varphi}(x, 0) + \tilde{\varphi}(-x, 0)$$

for all $x \in X$.

Proof. It follows from (2.8) that

$$\left\| g(x) - \frac{1}{4}g(2x) \right\| \leq \frac{1}{4}\varphi(2x, 0) + \frac{1}{4}\varphi(-2x, 0)$$

for all $x \in X$. So

$$(2.13) \quad \left\| \frac{1}{4^l}g(2^l x) - \frac{1}{4^m}g(2^m x) \right\| \leq \sum_{j=l+1}^m \frac{1}{4^j}\varphi(2^j x, 0) + \sum_{j=l+1}^m \frac{1}{4^j}\varphi(-2^j x, 0)$$

for all nonnegative integers m and l with $m > l$ and all $x \in X$. It follows from (2.11) and (2.13) that the sequence $\{\frac{1}{4^k}g(2^k x)\}$ is Cauchy for all $x \in X$. Since Y is complete, the sequence $\{\frac{1}{4^k}g(2^k x)\}$ converges. So one can define the mapping $Q : X \rightarrow Y$ by

$$Q(x) := \lim_{k \rightarrow \infty} \frac{1}{4^k}g(2^k x)$$

for all $x \in X$.

By (2.4) and (2.11),

$$\begin{aligned} \|DQ(x, y)\| &= \lim_{k \rightarrow \infty} \frac{1}{4^k} \|Dg(2^k x, 2^k y)\| \\ &\leq \lim_{k \rightarrow \infty} \frac{1}{4^k} (\varphi(2^k x, 2^k y) + \varphi(-2^k x, -2^k y)) = 0 \end{aligned}$$

for all $x, y \in X$. So $DQ(x, y) = 0$. Since $g : X \rightarrow Y$ is even, $Q : X \rightarrow Y$ is even. So the mapping $Q : X \rightarrow Y$ is Jensen quadratic. Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (2.13), we get (2.12). So there exists a Jensen quadratic mapping $Q : X \rightarrow Y$ satisfying (2.12).

The rest of the proof is similar to the proof of Theorem 2.2. $\square$

COROLLARY 2.5. *Let $p < 2$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping satisfying (2.10). Then there exists a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that*

$$\|f(x) + f(-x) - Q(x)\| \leq \frac{2^{p+1}\theta}{4 - 2^p} \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 2.4 to get the desired result. $\square$

3. Jensen additive mappings associated with inner product spaces

In this section, we prove the generalized Hyers-Ulam stability of the functional equation $Df(x, y) = 0$ in real Banach spaces: an odd case.

THEOREM 3.1. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ such that*

$$(3.1) \quad \Phi(x, y) := \sum_{j=0}^{\infty} 2^j \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}\right) < \infty,$$

$$(3.2) \quad \|Df(x, y)\| \leq \varphi(x, y)$$

for all $x, y \in X$. Then there exists a unique Jensen additive mapping $A : X \rightarrow Y$ such that

$$(3.3) \quad \|f(x) - f(-x) - A(x)\| \leq \Phi(x, 0) + \Phi(-x, 0)$$

for all $x \in X$.

Proof. Letting $y = 0$ in (3.2), we get

$$(3.4) \quad \left\| 3f\left(\frac{x}{2}\right) + f\left(\frac{-x}{2}\right) - f(x) \right\| \leq \varphi(x, 0)$$

for all $x \in X$. Replacing x by $-x$ in (3.4), we get

$$(3.5) \quad \left\| 3f\left(\frac{-x}{2}\right) + f\left(\frac{x}{2}\right) - f(-x) \right\| \leq \varphi(-x, 0)$$

for all $x \in X$. Let $h(x) := f(x) - f(-x)$ for all $x \in X$. It follows from (3.4) and (3.5) that

$$(3.6) \quad \left\| h(x) - 2h\left(\frac{x}{2}\right) \right\| \leq \varphi(x, 0) + \varphi(-x, 0)$$

for all $x \in X$. Hence

$$(3.7) \quad \begin{aligned} \|2^l h\left(\frac{x}{2^l}\right) - 2^m h\left(\frac{x}{2^m}\right)\| &\leq \sum_{j=l}^{m-1} 2^j \varphi\left(\frac{x}{2^j}, 0\right) \\ &\quad + \sum_{j=l}^{m-1} 2^j \varphi\left(-\frac{x}{2^j}, 0\right) \end{aligned}$$

for all nonnegative integers m and l with $m > l$ and all $x \in X$. It follows from (3.1) and (3.7) that the sequence $\{2^k h(\frac{x}{2^k})\}$ is Cauchy for

all $x \in X$. Since Y is complete, the sequence $\{2^k h(\frac{x}{2^k})\}$ converges. So one can define the mapping $A : X \rightarrow Y$ by

$$A(x) := \lim_{k \rightarrow \infty} 2^k h\left(\frac{x}{2^k}\right)$$

for all $x \in X$.

By (3.1) and (3.2),

$$\begin{aligned} \|DA(x, y)\| &= \lim_{k \rightarrow \infty} 2^k \left\| Dh\left(\frac{x}{2^k}, \frac{y}{2^k}\right) \right\| \\ &\leq \lim_{k \rightarrow \infty} 2^k \left(\varphi\left(\frac{x}{2^k}, \frac{y}{2^k}\right) + \varphi\left(-\frac{x}{2^k}, -\frac{y}{2^k}\right) \right) = 0 \end{aligned}$$

for all $x, y \in X$. So $DA(x, y) = 0$. Since $h : X \rightarrow Y$ is odd, $A : X \rightarrow Y$ is odd. So the mapping $A : X \rightarrow Y$ is Jensen additive. Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (3.7), we get (3.3). So there exists a Jensen additive mapping $A : X \rightarrow Y$ satisfying (3.3).

The rest of the proof is similar to the proof of Theorem 2.2. $\square$

COROLLARY 3.2. *Let $p > 1$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping such that*

$$(3.8) \quad \|Df(x, y)\| \leq \theta(\|x\|^p + \|y\|^p)$$

for all $x, y \in X$. Then there exists a unique Jensen additive mapping $A : X \rightarrow Y$ such that

$$\|f(x) - f(-x) - A(x)\| \leq \frac{2^{p+1}\theta}{2^p - 2} \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 3.1 to get the desired result. $\square$

THEOREM 3.3. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ satisfying (3.2) such that*

$$(3.9) \quad \Phi(x, y) := \sum_{j=1}^{\infty} 2^{-j} \varphi(2^j x, 2^j y) < \infty$$

for all $x, y \in X$. Then there exists a unique Jensen additive mapping $A : X \rightarrow Y$ such that

$$(3.10) \quad \|f(x) - f(-x) - A(x)\| \leq \Phi(x, 0) + \Phi(-x, 0)$$

for all $x \in X$.

Proof. It follows from (3.6) that

$$\left\| h(x) - \frac{1}{2}h(2x) \right\| \leq \frac{1}{2}\varphi(2x, 0) + \frac{1}{2}\varphi(-2x, 0)$$

for all $x \in X$. So

$$(3.11) \quad \left\| \frac{1}{2^l}h(2^l x) - \frac{1}{2^m}h(2^m x) \right\| \leq \sum_{j=l+1}^m \frac{1}{2^j}\varphi(2^j x, 0) + \sum_{j=l+1}^m \frac{1}{2^j}\varphi(-2^j x, 0)$$

for all nonnegative integers m and l with $m > l$ and all $x \in X$. It follows from (3.9) and (3.11) that the sequence $\{\frac{1}{2^k}h(2^k x)\}$ is Cauchy for all $x \in X$. Since Y is complete, the sequence $\{\frac{1}{2^k}h(2^k x)\}$ converges. So one can define the mapping $A : X \rightarrow Y$ by

$$A(x) := \lim_{k \rightarrow \infty} \frac{1}{2^k}h(2^k x)$$

for all $x \in X$.

By (3.2) and (3.9),

$$\begin{aligned} \|DA(x, y)\| &= \lim_{k \rightarrow \infty} \frac{1}{2^k} \|Dh(2^k x, 2^k y)\| \\ &\leq \lim_{k \rightarrow \infty} \frac{1}{2^k} (\varphi(2^k x, 2^k y) + \varphi(-2^k x, -2^k y)) = 0 \end{aligned}$$

for all $x, y \in X$. So $DA(x, y) = 0$. Since $h : X \rightarrow Y$ is odd, $A : X \rightarrow Y$ is odd. So the mapping $A : X \rightarrow Y$ is Jensen additive. Moreover, letting $l = 0$ and passing the limit $m \rightarrow \infty$ in (3.11), we get (3.10). So there exists a Jensen additive mapping $A : X \rightarrow Y$ satisfying (3.10).

The rest of the proof is similar to the proof of Theorem 2.2. $\square$

COROLLARY 3.4. *Let $p < 1$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping satisfying (3.8). Then there exists a unique Jensen additive mapping $A : X \rightarrow Y$ such that*

$$\|f(x) - f(-x) - A(x)\| \leq \frac{2^{p+1}\theta}{2-2^p} \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 3.3 to get the desired result. $\square$

Note that

$$\sum_{j=0}^{\infty} 2^j \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}\right) \leq \sum_{j=0}^{\infty} 4^j \varphi\left(\frac{x}{2^j}, \frac{y}{2^j}\right).$$

Combining Theorem 2.2 and Theorem 3.1, we obtain the following result.

THEOREM 3.5. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ satisfying (2.3) and (2.4). Then there exist a unique Jensen additive mapping $A : X \rightarrow Y$ and a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that*

$$\|2f(x) - A(x) - Q(x)\| \leq \tilde{\varphi}(x, 0) + \tilde{\varphi}(-x, 0) + \Phi(x, 0) + \Phi(-x, 0)$$

for all $x \in X$, where $\tilde{\varphi}$ and Φ are defined in (2.3) and (3.1), respectively.

COROLLARY 3.6. *Let $p > 2$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping satisfying (2.10). Then there exist a unique Jensen additive mapping $A : X \rightarrow Y$ and a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that*

$$\|2f(x) - A(x) - Q(x)\| \leq \left(\frac{2^{p+1}}{2^p - 2} + \frac{2^{p+1}}{2^p - 4} \right) \theta \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 3.5 to get the desired result. $\square$

Note that

$$\sum_{j=1}^{\infty} 4^{-j} \varphi(2^j x, 2^j y) \leq \sum_{j=1}^{\infty} 2^{-j} \varphi(2^j x, 2^j y).$$

Combining Theorem 2.4 and Theorem 3.3, we obtain the following result.

THEOREM 3.7. *Let $f : X \rightarrow Y$ be a mapping satisfying $f(0) = 0$ for which there exists a function $\varphi : X^2 \rightarrow [0, \infty)$ satisfying (2.4) and (3.9). Then there exist a unique Jensen additive mapping $A : X \rightarrow Y$ and a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that*

$$\|2f(x) - A(x) - Q(x)\| \leq \tilde{\varphi}(x, 0) + \tilde{\varphi}(-x, 0) + \Phi(x, 0) + \Phi(-x, 0)$$

for all $x \in X$, where $\tilde{\varphi}$ and Φ are defined in (2.11) and (3.9), respectively.

COROLLARY 3.8. *Let $p < 1$ and θ be positive real numbers, and let $f : X \rightarrow Y$ be a mapping satisfying (2.10). Then there exist a unique Jensen additive mapping $A : X \rightarrow Y$ and a unique Jensen quadratic mapping $Q : X \rightarrow Y$ such that*

$$\|2f(x) - A(x) - Q(x)\| \leq \left(\frac{2^{p+1}}{2-2^p} + \frac{2^{p+1}}{4-2^p} \right) \theta \|x\|^p$$

for all $x \in X$.

Proof. Define $\varphi(x, y) = \theta(\|x\|^p + \|y\|^p)$, and apply Theorem 3.7 to get the desired result. $\square$

References

- [1] T. Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan **2** (1950), 64–66.
- [2] P.W. Cholewa, *Remarks on the stability of functional equations*, Aequationes Math. **27** (1984), 76–86.
- [3] S. Czerwik, *On the stability of the quadratic mapping in normed spaces*, Abh. Math. Sem. Univ. Hamburg **62** (1992), 59–64.
- [4] P. Găvruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl. **184** (1994), 431–436.
- [5] D.H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [6] Th.M. Rassias, *On the stability of the linear mapping in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [7] Th.M. Rassias, *New characterizations of inner product spaces*, Bull. Sci. Math. **108** (1984), 95–99.
- [8] Th.M. Rassias, *On the stability of the quadratic functional equation and its applications*, Studia Univ. Babes-Bolyai **XLIII** (1998), 89–124.
- [9] Th.M. Rassias, *The problem of S.M. Ulam for approximately multiplicative mappings*, J. Math. Anal. Appl. **246** (2000), 352–378.
- [10] Th.M. Rassias, *On the stability of functional equations in Banach spaces*, J. Math. Anal. Appl. **251** (2000), 264–284.
- [11] Th.M. Rassias, *On the stability of functional equations and a problem of Ulam*, Acta Appl. Math. **62** (2000), 23–130.
- [12] Th.M. Rassias and P. Šemrl, *On the Hyers-Ulam stability of linear mappings*, J. Math. Anal. Appl. **173** (1993), 325–338.
- [13] Th.M. Rassias and K. Shibata, *Variational problem of some quadratic functionals in complex analysis*, J. Math. Anal. Appl. **228** (1998), 234–253.
- [14] F. Skof, *Proprietà locali e approssimazione di operatori*, Rend. Sem. Mat. Fis. Milano **53** (1983), 113–129.
- [15] S.M. Ulam, *Problems in Modern Mathematics*, Wiley, New York, 1960.

*

Department of Mathematics
Hanyang University
Seoul 133-791, Republic of Korea
E-mail: baak@hanyang.ac.kr

**

Seoul Science High School
Seoul 110-530, Republic of Korea
E-mail: gasop@naver.com

Seoul Science High School
Seoul 110-530, Republic of Korea
E-mail: lion328@naver.com

Seoul Science High School
Seoul 110-530, Republic of Korea
E-mail: bv2@naver.com

Seoul Science High School
Seoul 110-530, Republic of Korea
E-mail: sofyhsh92@naver.com