

THAINE'S THEOREM IN FUNCTION FIELD

HWANYUP JUNG*

ABSTRACT. Let F be a finite real abelian extension of a global function field k with $G = \text{Gal}(F/k)$. Assume that F is an extension field of the Hilbert class field K_ϵ of k and is contained in a cyclotomic function field K_n . Let ℓ be any prime number not dividing $ph_k|G|$. In this paper, we show that if $\theta \in \mathbb{Z}[G]$ annihilates the Sylow ℓ -subgroup of $\mathcal{O}_F^\times/\mathcal{C}_F$, then $(q-1)\theta$ annihilates the Sylow ℓ -subgroup of $\mathcal{C}l_F$.

1. Introduction

Let F be a totally real abelian number field with $G = \text{Gal}(F/\mathbb{Q})$. Let $\mathcal{O}_F^\times$ be the group of global units of F , $\mathcal{C}_F$ the group of cyclotomic units of F of conductor level and $\mathcal{C}l_F$ the class group of F . Let p be a prime not dividing $[F : \mathbb{Q}]$. In [4], Thaine has shown the following remarkable result: *If $\theta \in \mathbb{Z}[G]$ annihilates the Sylow p -subgroup of $\mathcal{O}_F^\times/\mathcal{C}_F$, then 2θ annihilates the Sylow p -subgroup of $\mathcal{C}l_F$.*

In this paper we consider the analogous problem in function fields. Let k be a global function field over the finite field $\mathbb{F}_q$ with q elements of characteristic p . Fix a place ∞ of k of degree one and a sign function $\text{sgn} : k_\infty \rightarrow \mathbb{F}_q$ with $\text{sgn}(0) = 0$, where k_∞ is the completion of k at ∞ . Let $\mathbb{A}$ be the Dedekind subring of k consisting of the functions regular away from ∞ . For any finite separable extension F of k , write $\mathcal{O}_F$ for the integral closure of $\mathbb{A}$ in F and $\mathcal{O}_F^\times$ be its group of units. Also write $\mathcal{C}l_F$ for the ideal class group of $\mathcal{O}_F$ and $h_F = |\mathcal{C}l_F|$. In this paper we only consider finite abelian extension F of k contained in a cyclotomic function field of (k, ∞, sgn) . Let F be a finite real abelian extension of k , i.e., ∞ splits completely in F , with $G = \text{Gal}(F/k)$. We assume that F contains the Hilbert class field of (k, ∞) . Let $\mathcal{C}_F$ be the group

Received November 28, 2008; Accepted February 13, 2009.

2000 Mathematics Subject Classification: Primary 11R58, 11R60, 11R18, 11R29.

Key words and phrases: class group, annihilator, function field.

This work was supported by the research grant of the Chungbuk National University in 2008.

of cyclotomic units of F in the sense of [1] (see the definition 2.2). The main result of this paper is

THEOREM 1.1. *Let ℓ be a prime not dividing $\ell \nmid ph_k|G|$. If $\theta \in \mathbb{Z}[G]$ annihilates the Sylow ℓ -subgroup of $\mathcal{O}_F^\times/\mathcal{C}_F$, then $(q-1)\theta$ annihilates the Sylow ℓ -subgroup of $\mathcal{C}l_F$.*

For any proper integral ideal of $\mathbb{A}$ of $\mathbf{n}$, we write $\Phi(\mathbf{n}) = |(\mathbb{A}/\mathbf{n})^\times|$ and denote by $K_{\mathbf{n}}$ the cyclotomic function field of (k, ∞, sgn) of conductor $\mathbf{n}$. Moreover we write $K_{\mathfrak{e}}$ for the Hilbert class field of (k, ∞) . For details in the theory of sgn-normalized Drinfeld $\mathbb{A}$ -module and cyclotomic function field over the global function field k , we refer the readers to [2] or [3].

2. Preliminary

2.1. Cyclotomic units

Let ρ be a fixed sgn-normalized Drinfeld $\mathbb{A}$ -module. For any proper ideal $\mathbf{n}$ of $\mathbb{A}$, we fix a primitive $\mathbf{n}$ -torsion point $\lambda_{\mathbf{n}}$ of ρ .

LEMMA 2.1. *Let $\mathbf{n} = \mathfrak{p}\mathfrak{f}$, where $\mathfrak{f} \neq \mathfrak{e}$ and $\mathfrak{p}$ is a prime ideal of $\mathbb{A}$. Then we have*

$$N_{K_{\mathbf{n}}/K_{\mathfrak{f}}}(\lambda_{\mathbf{n}}) = \begin{cases} \rho_{\mathfrak{p}}(\lambda_{\mathbf{n}}), & \text{if } \mathfrak{p}|\mathfrak{f}, \\ \rho_{\mathfrak{p}}(\lambda_{\mathbf{n}})^{1-\sigma_{\mathfrak{p}}^{-1}}, & \text{if } \mathfrak{p} \nmid \mathfrak{f}, \end{cases}$$

where $\sigma_{\mathfrak{p}}$ denotes the Frobenius automorphism of $\mathfrak{p}$ in $K_{\mathfrak{f}}$.

Let F be a finite real abelian extension of k . Let $\mathbf{m}$ be the conductor of F , i.e. $K_{\mathbf{m}}$ is the smallest cyclotomic function field containing F . For each proper ideal $\mathfrak{f}$ of $\mathbb{A}$, put $F_{\mathfrak{f}} = F \cap K_{\mathfrak{f}}$ and $\lambda_{\mathfrak{f},F} = N_{K_{\mathfrak{f}}/F_{\mathfrak{f}}}(\lambda_{\mathfrak{f}})$.

DEFINITION 2.2. Let $\mathcal{D}_F$ be the G -submodule of $F^\times$ generated by $\mathbb{F}_q^\times$ and $\lambda_{\mathfrak{f},F}$ with all $\mathfrak{f} \neq \mathfrak{e}$. We define $\mathcal{C}_F := \mathcal{D}_F \cap \mathcal{O}_F^\times$, called the *group of cyclotomic units* of F .

As a G -module, $\mathcal{D}_F$ is generated by $\mathbb{F}_q^\times \cup \{\lambda_{\mathfrak{f},F} : \mathfrak{e} \neq \mathfrak{f}|\mathbf{m}\} \cup \{N_{K_{\mathfrak{p}}/K_{\mathfrak{e}}}(\lambda_{\mathfrak{p}})\}_{\mathfrak{p}}$, where $\mathfrak{p}$ runs over all prime ideals of $\mathbb{A}$ such that $\mathfrak{p} \nmid \mathbf{m}$.

For any prime ideal $\mathfrak{q}$ of $\mathbb{A}$ which splits completely in F , write $F(\mathfrak{q}) = F \cdot K_{\mathfrak{q}}$. Since all prime ideals of $\mathcal{O}_F$ above $\mathfrak{q}$ are totally ramified in $F(\mathfrak{q})$, G acts on the set of prime ideals of $\mathcal{O}_{F(\mathfrak{q})}$ above $\mathfrak{q}$.

PROPOSITION 2.3. *For any unit $\varepsilon \in \mathcal{C}_F$, there exists $u \in \mathcal{O}_{F(\mathfrak{q})}^\times$ such that $N_{F(\mathfrak{q})/F}(u) = 1$ and $u \equiv \varepsilon \pmod{\tilde{\mathfrak{Q}}^\sigma}$ for all $\sigma \in G$.*

Proof. It suffices to show that for any $\varepsilon \in \mathbb{F}_q^\times \cup \{\lambda_{f,F} : \mathfrak{e} \neq f|\mathfrak{m}\} \cup \{N_{K_p/K_e}(\lambda_p)\}_{\mathfrak{p}}$, where $\mathfrak{p}$ runs over all prime ideals of $\mathbb{A}$ such that $\mathfrak{p} \nmid \mathfrak{m}$ and $\mathfrak{p} \neq \mathfrak{q}$, there exists $u \in \mathcal{O}_{F(\mathfrak{q})}^\times$ satisfying the required conditions. For $\varepsilon \in \mathbb{F}_q^\times$, it is easy to see that $u = \varepsilon$ satisfies the required ones. For $\varepsilon = \lambda_{f,F}$ with $\mathfrak{e} \neq f|\mathfrak{m}$, take $u = \prod_{\tau \in \text{Gal}(K_f/F_f)} (\lambda_f^\tau + \lambda_q)$. Then $u \in F_f(\mathfrak{q}) \subset F(\mathfrak{q})$, and so $u \in \mathcal{O}_{F(\mathfrak{q})}^\times$. Since $N_{F(\mathfrak{q})/F}(u) = N_{K_{fq}/K_f}(u)$, we have

$$N_{F(\mathfrak{q})/F}(u) = \prod_{\tau \in \text{Gal}(K_f/F_f)} \left(\frac{\rho_q(\lambda_f^\tau)}{\lambda_f^\tau} \right) = N_{K_f/F_f}(\lambda_f)^{\sigma_q-1} = 1.$$

Since λ_q is contained in $\tilde{\mathfrak{Q}}^\sigma$ for all $\sigma \in G$, $u \equiv \prod_{\tau \in \text{Gal}(K_f/F_f)} \lambda_f^\tau = \lambda_{f,F} \pmod{\tilde{\mathfrak{Q}}^\sigma}$. For $\varepsilon = N_{K_p/K_e}(\lambda_p)$ with $\mathfrak{p} \nmid \mathfrak{m}, \mathfrak{p} \neq \mathfrak{q}$, similar as above, we can show that $u = \prod_{\tau \in \text{Gal}(K_p/K_e)} (\lambda_p^\tau + \lambda_q)$ satisfies the required ones. $\square$

Fix a generator s of $(\mathbb{A}/\mathfrak{q})^\times$ and let $\tau \in \text{Gal}(F(\mathfrak{q})/F)$ be the automorphism such that $\tau(\lambda_q) = \rho_s(\lambda_q)$. Let u be a unit in Lemma 2.3 and choose $w \in F(\mathfrak{q})$ satisfying $w^\tau = uw$. Then we have $(w) = \mathfrak{D} \prod_{\sigma \in G} (\tilde{\mathfrak{Q}}^{\sigma^{-1}})^{r_\sigma}$, where $\mathfrak{D}$ is the lift of an ideal of $\mathcal{O}_F$ relatively prime to $\mathfrak{q}$ and $r_\sigma \in \mathbb{Z}$. As in classical case, r_σ is determined uniquely modulo $\Phi(\mathfrak{q})$ by $s^{r_\sigma} \equiv \sigma(\varepsilon) \pmod{\mathfrak{Q}}$.

2.2. Applications of Tchebotarev density theorem

Given ideal class $\mathfrak{c}$ of $\mathcal{O}_F$ and a positive integer N , we define $\mathbf{P}(\mathfrak{c}, N)$ as the set of prime ideals $\mathfrak{Q}$ belonging to $\mathfrak{c}$ and lying above a prime ideal $\mathfrak{q}$ of $\mathbb{A}$ which splits completely in F and $\Phi(\mathfrak{q}) \equiv 0 \pmod{N}$. By using the Tchebotarev density theorem ([3, Theorem 9.13 A]), we can show that if N is a positive integer with $p \nmid N$, then $\mathbf{P}(\mathfrak{c}, N)$ is an infinite set.

LEMMA 2.4. *For any nonconstant $z \in k_\infty^\times$ and a positive integer c prime to p , $k_\infty(\sqrt[c]{z})$ is a totally ramified extension over k_∞ .*

Proof. It is an easy consequence of local class field theory. $\square$

PROPOSITION 2.5. *Let N and c be positive integers with $c|N$ and $p \nmid N$. Let x be a nonconstant element of $\mathcal{O}_F$. Suppose that for all (except possibly a finite set) of the prime ideals $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, N)$, there exists $y_{\mathfrak{Q}} \in \mathcal{O}_F$ such that $x \equiv y_{\mathfrak{Q}}^c \pmod{\mathfrak{Q}}$. Then $x = \alpha y^{c/f}$ for some $y \in \mathcal{O}_F$ and $\alpha \in \mathbb{F}_q^*$, a f -th root of unity, where $f = \gcd(c, q-1)$.*

Proof. Fix $\zeta_c \in \overline{\mathbb{F}}_q$ a primitive c -th root of unity and let $\sqrt[c]{x}$ be a fixed c -th root of x . By Lemma 2.4, $F(\sqrt[c]{x})/F$ is a geometric extension and $\tilde{F} \cap F(\sqrt[c]{x}) = F$, where $\tilde{F} = F(\zeta_c)$. Let L be the Galois closure of $F(\sqrt[c]{x})$ over F . Clearly, $L \subseteq \tilde{F}(\sqrt[c]{x})$. Let $F' = \mathbb{F}_{q^d}F \subseteq \tilde{F}$, where $d = [L : F(\sqrt[c]{x})]$. Since $F(\sqrt[c]{x})/F$ is a geometric extension, $L = F'F(\sqrt[c]{x}) = F'(\sqrt[c]{x})$, $\tilde{F} \cap L = F'$ and $F' \cap F(\sqrt[c]{x}) = F$. Let $f = \gcd(c, q^d - 1)$ and $\zeta_f = (\zeta_c)^{c/f}$. Clearly, $F(\zeta_f) \subseteq F'$. Let $Q(X)$ be the irreducible polynomial of $\sqrt[c]{x}$ over F . Then $Q(X) = \prod_{j \in J} (X - \zeta_c^j \sqrt[c]{x})$, where J is a subset of $\{1, 2, \dots, c\}$ and $\zeta_c^j \in \mathbb{F}_{q^d}$ for all $j \in J$. It is easy to see that $\zeta_c^j \in \mathbb{F}_{q^d}$ if and only if j divides c/f . Hence $Q(X) = \prod_{j \in J^*} (X - \zeta_f^j \sqrt[c]{x})$ for some $J^* \subseteq \{1, 2, \dots, f\}$. Since $L = F'F(\sqrt[c]{x})$ and $F' \cap F(\sqrt[c]{x}) = F$, $Q(X)$ is also the irreducible polynomial of $\sqrt[c]{x}$ over F' . Since $L = F(\sqrt[c]{x}, \zeta_f^j : j \in J^*) \subseteq F(\sqrt[c]{x}, \zeta_f) \subseteq F'(\sqrt[c]{x}) = L$, we have $L = F(\sqrt[c]{x}, \zeta_f)$ and $F' = F(\zeta_f)$. Note that L/F' is a finite Galois extension and let $\Gamma = \text{Gal}(L/F')$. Let $\mathcal{U}_f$ be the group of f -th roots of unity in $\overline{\mathbb{F}}_q$. Since $Q(X) = \prod_{\sigma \in \Gamma} (X - \sigma(\sqrt[c]{x}))$, $(\sqrt[c]{x})^{\sigma^{-1}} \in \mathcal{U}_f$ for all $\sigma \in \Gamma$. Let $\psi : \Gamma \rightarrow \mathcal{U}_f$ be the homomorphism defined by $\psi(\sigma) = (\sqrt[c]{x})^{\sigma^{-1}}$. Then $\text{Im}(\psi) = \langle \zeta_{f_0} \rangle$ for some $f_0 | f$, and so $Q(X) = \prod_{j=1}^{f_0} (X - \zeta_{f_0}^j \sqrt[c]{x})$. Now we can follow the same argument as in the classical case for the rest of proof. $\square$

For each nonconstant unit $x \in \mathcal{O}_F^\times$, we define the number $\phi(x)$ as the greatest positive integer n such that $x = u^n$ for some $u \in F$. Clearly, $\phi(\sigma(x)) = \phi(x)$ for all $\sigma \in G$. Fix a triple $(x, \mathfrak{c}, N)$, where $x \in \mathcal{C}_F$, $\mathfrak{c}$ an ideal class and N a positive integer with $p \nmid N$. For each $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, N)$, let $s_{\mathfrak{Q}}$ be a fixed generator of $(\mathbb{A}/\mathfrak{q})^\times$, where $\mathfrak{q} = \mathfrak{Q} \cap \mathbb{A}$. Then there exists a nonzero fractional ideal $\mathfrak{D}_{\mathfrak{Q}}$ of $\mathcal{O}_F$ such that $\mathfrak{D}_{\mathfrak{Q}}^N \prod_{\sigma \in G} (\mathfrak{Q}^{\sigma^{-1}})^{r_{\sigma}(\mathfrak{Q})}$ is a principal ideal, where the integers $r_{\sigma}(\mathfrak{Q})$ satisfy $s_{\mathfrak{Q}}^{r_{\sigma}(\mathfrak{Q})} \equiv \sigma(x) \pmod{\mathfrak{Q}}$. Let $\sigma \in G$ be fixed. We define $g = g(x, \mathfrak{c}, N, \sigma)$ as the greatest common divisor of N and of all the $r_{\sigma}(\mathfrak{Q})$ such that $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, N)$.

Fix an embedding of F into k_∞ . We call $x \in F$ *positive* if $\text{sgn}(x) = 1$.

THEOREM 2.6. *Given $x \in \mathcal{C}_F \setminus \mathbb{F}_q^\times$, $\mathfrak{c}$ an ideal class of $\mathcal{O}_F$, N a positive integer with $p \nmid N$ and $\sigma \in G$, let $g = g(x, \mathfrak{c}, N, \sigma)$. If $\gcd(N, q-1) = 1$ or $\sigma(x)$ is a positive, then we have $\gcd(N, \phi(x)) | g | \gcd(N, f\phi(x))$, where $f = \gcd(g, q-1)$.*

Proof. As in classical case, it can be shown easily that $\gcd(N, \phi(x))$ divides g . For any $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, N)$, there exists $y_{\mathfrak{Q}} \in \mathbb{A}$ such that $\sigma(x) \equiv y_{\mathfrak{Q}}^g \pmod{\mathfrak{Q}}$. By Proposition 2.5, $\sigma(x) = \alpha y^{g/f}$ for some $y \in \mathcal{O}_F$ and $\alpha \in \mathbb{F}_q^\times$, a f -th root of unity, where $f = \gcd(g, q-1)$. If $\gcd(N, q-1) = 1$,

then $f = 1, \alpha = 1$ and hence $\sigma(x) = y^{g/f}$. If $\sigma(x)$ is a positive, then $1 = \alpha \cdot \text{sgn}(y)^{g/f}$. Replacing y by $y/\text{sgn}(y)$, we also have $\sigma(x) = y^{g/f}$. Thus $g|f\phi(x)$, and so $g|(N, f\phi(x))$. $\square$

3. Proof of Theorem 1.1

For any finite abelian group A , we denote by A_ℓ the Sylow ℓ -subgroup of A for any prime ℓ . Let $x \in \mathcal{C}_F, \mathfrak{c} \in \mathcal{C}l_F$ and N positive integer with $p \nmid N$ be given. For any $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, N)$, there is an ideal class $\mathfrak{d}_{\mathfrak{Q}} \in \mathcal{C}l_F$ such that

$$(3.1) \quad \mathfrak{d}_{\mathfrak{Q}}^N \prod_{\sigma \in G} \sigma^{-1}(\mathfrak{c})^{r_{\sigma}(\mathfrak{Q})} = 1,$$

where the integers $r_{\sigma}(\mathfrak{Q})$ satisfy $s_{\mathfrak{Q}}^{r_{\sigma}(\mathfrak{Q})} \equiv \sigma(x) \pmod{\mathfrak{Q}}$.

PROPOSITION 3.1. *Let ℓ be a prime number such that $\ell \neq p$. Let $x \in \mathcal{C}_F$ and let ℓ^n be an exponent of $(\mathcal{C}l_F)_{\ell}$. If $\mathfrak{c} \in (\mathcal{C}l_F)_{\ell}, \mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, \ell^n)$ and if $r_{\sigma} = r_{\sigma}(\mathfrak{Q}), \sigma \in G$, are integers satisfying $s_{\mathfrak{Q}}^{r_{\sigma}} \equiv \sigma(x) \pmod{\mathfrak{Q}}$, then $\varrho = \varrho_{\mathfrak{Q}} = \sum_{\sigma \in G} r_{\sigma}(\mathfrak{Q})\sigma^{-1}$ annihilates $\mathfrak{c}$.*

Proof. Note that the integers r_{σ} are uniquely determined modulo ℓ^n because $\ell^n | \Phi(\mathfrak{q})$. Since $\mathfrak{c}^{\ell^n} = 1$, (3.1) holds with $N = \ell^n$. Since all conjugates of $\mathfrak{c}$ belong to $(\mathcal{C}l_F)_{\ell}$, $\mathfrak{d}_{\mathfrak{Q}}^{\ell^n} \in (\mathcal{C}l_F)_{\ell}$, and hence $\mathfrak{d}_{\mathfrak{Q}} \in (\mathcal{C}l_F)_{\ell}$ and $\mathfrak{d}_{\mathfrak{Q}}^{\ell^n} = 1$. Therefore $\mathfrak{c}^{\varrho} = 1$. $\square$

For any character χ of G , we define the idempotent element e_{χ} as follows:

$$e_{\chi} = \frac{1}{|G|} \sum_{\sigma \in G} \text{Tr}(\chi(\sigma))\sigma^{-1} \in \mathbb{Q}_{\ell}[G],$$

where "Tr" is the trace map from $\mathbb{Q}_{\ell}(\chi) := \mathbb{Q}_{\ell}(\chi(\sigma) : \sigma \in G)$ to $\mathbb{Q}_{\ell}$. Now assume that $|G|$ is prime to ℓ . Then $e_{\chi} \in \mathbb{Z}_{\ell}[G]$ for all character χ of G .

Since there is a Minkowski unit in $\mathcal{O}_F^{\times}$, as in classical case, we have

PROPOSITION 3.2. *Suppose that $\ell \nmid p|G|$. Let χ be a nontrivial character of G and ℓ^n an exponent of $(\mathcal{O}_F^{\times}/\mathcal{C}_F)_{\ell}$. Let $\ell^{a_{\chi}}$ be the exact exponent of the χ -part $e_{\chi}(\mathcal{O}_F^{\times}/\mathcal{C}_F)_{\ell}$ of $(\mathcal{O}_F^{\times}/\mathcal{C}_F)_{\ell}$. Then there exists a positive $x \in e_{\chi}(\mathcal{C}_F/\mathcal{C}_F \cap (\mathcal{O}_F^{\times})^{\ell^n})$ such that $\ell^{a_{\chi}}|\phi(x)$.*

LEMMA 3.3. *Suppose $\theta \in e_{\chi}\mathbb{Z}/\ell^n\mathbb{Z}[G]$ and ℓ^a is the highest power of ℓ dividing θ ($0 \leq a < n$). Then $\ell^{-a}\theta e_{\chi}\mathbb{Z}/\ell^n\mathbb{Z}[G] = e_{\chi}\mathbb{Z}/\ell^n\mathbb{Z}[G]$. In particular, there exists $\theta' \in e_{\chi}\mathbb{Z}/\ell^n\mathbb{Z}[G]$ such that $\ell^{-a}\theta\theta' = e_{\chi}$.*

Proof. See the proof of Lemma 15.6 in [5]. $\square$

Now we give the proof of Theorem 1.1. Let ℓ^n be an exponent of $(\mathcal{C}l_F)_\ell$ and $(\mathcal{O}_F^\times/\mathcal{C}_F)_\ell$. Let $\mathfrak{c} \in (\mathcal{C}l_F)_\ell$. For each $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, \ell^n)$, choose a generator $s_{\mathfrak{Q}}$ of $(\mathbb{A}/\mathfrak{q})^\times$, where $\mathfrak{q} = \mathfrak{Q} \cap \mathbb{A}$, and define

$$\Psi_{\mathfrak{Q}} : \mathcal{C}_F/\mathcal{C}_F \cap (\mathcal{O}_F^\times)^{\ell^n} \rightarrow \mathbb{Z}/\ell^n\mathbb{Z}[G]$$

by $\Psi_{\mathfrak{Q}}(x) = \sum_{\sigma \in G} r_\sigma \sigma^{-1}$, where the integers $r_\sigma = r_\sigma(\mathfrak{Q})$ are uniquely determined modulo ℓ^n and satisfy $s_{\mathfrak{Q}}^{r_\sigma} \equiv \sigma(x) \pmod{\mathfrak{Q}}$. By Proposition 3.1, $\mathfrak{c}^{\Psi_{\mathfrak{Q}}(x)} = 1$ for all $x \in \mathcal{C}_F/\mathcal{C}_F \cap (\mathcal{O}_F^\times)^{\ell^n}$. For any nontrivial character χ of G , let $\Psi_{\mathfrak{Q}}^\chi : e_\chi(\mathcal{C}_F/\mathcal{C}_F \cap (\mathcal{O}_F^\times)^{\ell^n}) \rightarrow e_\chi\mathbb{Z}/\ell^n\mathbb{Z}[G]$ be the restriction of $\Psi_{\mathfrak{Q}}$. Let ℓ^{a_χ} be the exact exponent of $e_\chi(\mathcal{O}_F^\times/\mathcal{C}_F)_\ell$. There exists a positive $x \in e_\chi(\mathcal{C}_F/\mathcal{C}_F \cap (\mathcal{O}_F^\times)^{\ell^n})$ satisfying $\ell^{a_\chi} \nmid \phi(x)$. If $\ell^e \mid (q-1)$, then $g = g(x, \mathfrak{c}, \ell^n, 1)$ divides $\ell^{a_\chi+e}$. There exists $\mathfrak{Q} \in \mathbf{P}(\mathfrak{c}, \ell^n)$ satisfying $\Psi_{\mathfrak{Q}}^\chi(x) \not\equiv 0 \pmod{\ell^{a_\chi+e+1}}$. For $\mathfrak{Q}$ as above, let a' be the minimal such that $\Psi_{\mathfrak{Q}}^\chi(x) \not\equiv 0 \pmod{\ell^{a'+1}}$, so that $a' \leq a_\chi + e$. Then $\ell^{-a'} \Psi_{\mathfrak{Q}}^\chi(x) \mathbb{Z}/\ell^n\mathbb{Z}[G] = e_\chi\mathbb{Z}/\ell^n\mathbb{Z}[G]$. Thus $\text{Im}(\Psi_{\mathfrak{Q}}^\chi) \supseteq \ell^{a'} e_\chi\mathbb{Z}/\ell^n\mathbb{Z}[G] \supseteq (q-1)\ell^{a_\chi} e_\chi\mathbb{Z}/\ell^n\mathbb{Z}[G]$, and so $(q-1)\ell^{a_\chi} e_\chi$ annihilates $\mathfrak{c}$. Since $\mathfrak{c} \in (\mathcal{C}l_F)_\ell$ is arbitrary, it proves that $(q-1)\ell^{a_\chi} e_\chi$ annihilates $(\mathcal{C}l_F)_\ell$. This is also true for the trivial character χ_0 because e_{χ_0} is essentially the norm and $\ell \nmid h_k$.

Now, suppose $\theta \in \mathbb{Z}[G]$ annihilates $(\mathcal{O}_F^\times/\mathcal{C}_F)_\ell$. For any character χ , θe_χ annihilates $e_\chi(\mathcal{O}_F^\times/\mathcal{C}_F)_\ell$. Let ℓ^b be the maximal power of ℓ dividing θe_χ . There exists $\theta' \in e_\chi\mathbb{Z}/\ell^n\mathbb{Z}[G]$ such that $\theta e_\chi \theta' = \ell^b e_\chi$ (by Lemma 3.3). Then ℓ^b annihilates $e_\chi(\mathcal{O}_F^\times/\mathcal{C}_F)_\ell$, and so $b \geq a_\chi$. Thus, $\ell^{a_\chi} \mid \theta e_\chi$ and $(q-1)\theta e_\chi$ annihilates $(\mathcal{C}l_F)_\ell$. Since $(q-1)\theta = \sum_\chi (q-1)\theta e_\chi$, it follows that $(q-1)\theta$ annihilates $(\mathcal{C}l_F)_\ell$.

References

- [1] J. Ahn, S. Bae and H. Jung, *Cyclotomic units and Stickelberger ideals of global function fields*. Trans. Amer. Math. Soc. **355** (2003), 1803-1818.
- [2] D. Hayes, *A brief introduction to Drinfeld modules*. The arithmetic of function fields (Columbus, OH, 1991), 1-32.
- [3] M. Rosen, *Number theory in function fields*. Graduate Texts in Mathematics, 210. Springer-Verlag, New York, 2002.
- [4] F. Thaine, *On the ideal class groups of real abelian number fields*. Ann. of Math. (2) **128** (1988), 1-18.
- [5] L. Washington, *Introduction to cyclotomic fields*. Second edition. GTM 83. Springer-Verlag, New York, (1997).

*

Department of Mathematics Education
Chungbuk National University
Cheongju 361-763, Republic of Korea
E-mail: `hyjung@chungbuk.ac.kr`