

ON WEAK M -SEMICONINUITY ON SPACES WITH MINIMAL STRUCTURES

WON KEUN MIN* AND YOUNG KEY KIM**

ABSTRACT. We introduce the notion of weak M -semicontinuity which is a generalization of M -semicontinuity defined between spaces with minimal structures. We also investigate some properties and characterizations for such a notion.

1. Introduction

Popa and Noiri [5] introduced the notion of minimal structure which is a generalization of a topology on a given nonempty set. And they introduced the notion of M -continuous function as a function defined between spaces with minimal structures. They showed that the M -continuous functions have properties similar to those of continuous functions between topological spaces.

In [2], we introduced the notions of m -semiopen sets, m -semi-interior and m -semi-closure operators on a space with a minimal structure. For a given nonempty set, if a minimal structure is a topology, the m -semiopen sets are semiopen sets defined in [1]. We also introduced the notion of M -semicontinuous function and studied characterizations for the M -semicontinuity in [2]. In this paper, we introduce the notion of weakly M -semicontinuous function which is a generalization of M -semicontinuous function defined between spaces with minimal structures. And we investigate characterizations for such a notion and relationship between weakly M -semicontinuous function and strongly M -semiclosed graph.

Received December 24, 2009; Accepted April 23, 2010.

2010 Mathematics Subject Classification: Primary 54C08.

Key words and phrases: weakly M -semicontinuous, M -semicontinuous, M -continuous, strongly M -semiclosed graph, m -Urysohn, m -semi- T_2 .

Correspondence should be addressed to Won Keun Min, wkmin@kangwon.ac.kr.

2. Preliminaries

A subfamily m_X of the power set $P(X)$ of a nonempty set X is called a *minimal structure* [5] on X if $\emptyset \in m_X$ and $X \in m_X$. By (X, m_X) , we denote a nonempty set X with a minimal structure m_X on X . Simply we call (X, m_X) a space with a minimal structure m_X on X . Let (X, m_X) be a space with a minimal structure m_X on X . For a subset A of X , the closure of A and the interior of A are defined as the following:

$$\begin{aligned} mInt(A) &= \cup\{U : U \subseteq A, U \in m_X\}; \\ mCl(A) &= \cap\{F : A \subseteq F, X - F \in m_X\}. \end{aligned}$$

THEOREM 2.1. [5] *Let (X, m_X) be a space with a minimal structure m_X on X and $A \subseteq X$.*

- (1) $X = mInt(X)$ and $\emptyset = mCl(\emptyset)$.
- (2) $mInt(A) \subseteq A$ and $A \subseteq mCl(A)$.
- (3) If $A \in m_X$, then $mInt(A) = A$ and if $X - F \in m_X$, then $mCl(F) = F$.
- (4) If $A \subseteq B$, then $mInt(A) \subseteq mInt(B)$ and $mCl(A) \subseteq mCl(B)$.
- (5) $mInt(mInt(A)) = mInt(A)$ and $mCl(mCl(A)) = mCl(A)$.
- (6) $mCl(X - A) = X - mInt(A)$ and $mInt(X - A) = X - mCl(A)$.

Let (X, m_X) be a space with a minimal structure m_X on X and $A \subset X$. A subset A of X is called an *m-semiopen set* [2] if $A \subseteq mCl(mInt(A))$. The complement of an *m-semiopen set* is called an *m-semiclosed set*. Let (X, m_X) be a space with a minimal structure m_X on X . For a subset A of X , the *m-semi-closure* of A and the *m-semi-interior* of A , denoted by $msCl(A)$ and $msInt(A)$, respectively, are defined as the following:

$$msCl(A) = \cap\{F : A \subseteq F, F \text{ is } m\text{-semiclosed in } X\};$$

$$msInt(A) = \cup\{U : U \subseteq A, U \text{ is } m\text{-semiopen in } X\}.$$

THEOREM 2.2. [2] *Let (X, m_X) be a space with a minimal structure m_X on X and $A \subseteq X$. Then*

- (1) $msInt(A) \subseteq A$ and $A \subseteq msCl(A)$.
- (2) If $A \subseteq B$, then $msInt(A) \subseteq msInt(B)$ and $msCl(A) \subseteq msCl(B)$.
- (3) A is *m-semiopen* iff $msInt(A) = A$.
- (4) F is *m-semiclosed* iff $msCl(F) = F$.
- (5) $msInt(msInt(A)) = msInt(A)$ and $msCl(msCl(A)) = msCl(A)$.
- (6) $msCl(X - A) = X - msInt(A)$ and $msInt(X - A) = X - msCl(A)$.

Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function between two spaces with minimal structures m_X and m_Y . Then f is said to be M -continuous [4] (resp. M -semicontinuous [3]) if for each x and each m -open set V containing $f(x)$, there exists an m -open (resp. m -semiopen) set U containing x such that $f(U) \subseteq V$.

3. Main Results

DEFINITION 3.1. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function between two spaces with minimal structures m_X and m_Y , respectively. Then f is said to be *weakly M -semicontinuous* if for each m -open set A of Y , $f^{-1}(A) \subseteq msInt(f^{-1}(msCl(A)))$.

Every M -semicontinuous function is weakly M -semicontinuous but the converse is not always true.

EXAMPLE 3.2. Let $X = Y = \{a, b, c\}$ and let us consider two minimal structures $m_X = \{\emptyset, \{a\}, \{b\}, X\}$ and $m_Y = \{\emptyset, \{a, b\}, \{b, c\}, Y\}$. Consider a function $f : (X, m_X) \rightarrow (Y, m_Y)$ defined as $f(a) = f(b) = a$, $f(c) = c$. Then for $\{a, b\}, \{b, c\} \in m_Y$, $msCl(\{a, b\}) = msCl(\{b, c\}) = Y$ in (Y, m_Y) , and so clearly f is weakly M -semicontinuous. But for $\{b, c\} \in m_Y$, $f^{-1}(\{b, c\})$ is not m -semiopen in (X, m_X) . Consequently f is not M -semicontinuous.

M -continuous \Rightarrow M -semicontinuous \Rightarrow weakly M -semicontinuous

THEOREM 3.3. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y . Then f is weakly M -semicontinuous if and only if for every point x and each m -open set V containing $f(x)$, there exists an m -semiopen set U containing x such that $f(U) \subseteq msCl(V)$.

Proof. Suppose f is a weakly M -semicontinuous function. Let x be in X and V an m -open set containing $f(x)$. Then there exists an m -semiopen set B such that $f(x) \in B \subseteq V$. From the hypothesis, it follows

$$x \in f^{-1}(B) \subseteq msInt(f^{-1}(msCl(B))) \subseteq msInt(f^{-1}(msCl(V))).$$

Since $x \in msInt(f^{-1}(msCl(V)))$, there exists an m -semiopen set U such that $x \in U \subseteq f^{-1}(msCl(V))$, and hence $f(U) \subseteq msCl(V)$.

For the converse, let V be an m -open set in Y . For each $x \in f^{-1}(V)$, there exists an m -semiopen set U containing x such that $f(U) \subseteq msCl(V)$. This implies

$$f^{-1}(V) \subseteq \cup\{U : x \in f^{-1}(V)\} \subseteq f^{-1}(msCl(V)).$$

Since $\cup\{U : x \in f^{-1}(V)\}$ is an m -semiopen set containing $f^{-1}(V)$, we have $f^{-1}(V) \subseteq msInt(f^{-1}(msCl(V)))$. Hence f is weakly M -semicontinuous. \square

A minimal structure m_X on a nonempty set X is said to have property (\mathcal{B}) [5] if the union of any family of subsets belonging to m_X belongs to m_X .

LEMMA 3.4. [5] *Let m_X be a minimal structure on a nonempty set X satisfying (\mathcal{B}) . For $A \subseteq X$, the following are equivalent:*

- (1) $A \in m_X$ if and only if $mInt(A) = A$.
- (2) A is m -closed if and only if $mCl(A) = A$.

THEOREM 3.5. *Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y , respectively. If m_Y has the property (\mathcal{B}) , then the following statements are equivalent:*

- (1) f is weakly M -semicontinuous.
- (2) $msCl(f^{-1}(msInt(F))) \subseteq f^{-1}(F)$ for each m -closed set F in Y .
- (3) $msCl(f^{-1}(V)) \subseteq f^{-1}(mCl(V))$ for each m -open set V in Y .

Proof. (1) \Rightarrow (2) Let F be any m -closed set of Y . Then from Theorem 2.2, it follows

$$\begin{aligned} f^{-1}(Y - F) &\subseteq msInt(f^{-1}(msCl(Y - F))) \\ &= msInt(f^{-1}(Y - msInt(F))) \\ &= msInt(Y - f^{-1}(msInt(F))) \\ &= X - msCl(f^{-1}(msInt(F))). \end{aligned}$$

Hence we have $msCl(f^{-1}(msInt(F))) \subseteq f^{-1}(F)$.

Similarly we can prove that the implication (2) \Rightarrow (1) is hold.

(2) \Rightarrow (3) For any m -open set V in Y , by Lemma 3.4, $mCl(V)$ is m -closed. Since $V \subseteq msInt(mCl(V))$, we have the following relation:

$$msCl(f^{-1}(V)) \subseteq msCl(f^{-1}(msInt(mCl(V)))) \subseteq f^{-1}(mCl(V)).$$

Hence the statement (3) is obtained.

(3) \Rightarrow (1) For each an m -open set V in Y . Then from

$$V \subseteq mInt(msCl(V)),$$

it follows

$$\begin{aligned}
 f^{-1}(V) &\subseteq f^{-1}(mInt(msCl(V))) \\
 &= X - f^{-1}(mCl(Y - msCl(V))) \\
 &\subseteq X - msCl(f^{-1}(Y - msCl(V))) \\
 &= msInt(f^{-1}(msCl(V))).
 \end{aligned}$$

Hence f is weakly M -semicontinuous. \square

THEOREM 3.6. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y . If m_Y has the property (\mathcal{B}) , then the following statements are equivalent:

- (1) f is weakly M -semicontinuous.
- (2) $msCl(f^{-1}(msInt(B))) \subseteq f^{-1}(mCl(B))$ for each set B in Y .
- (3) $f^{-1}(mInt(B)) \subseteq msInt(f^{-1}(msCl(B)))$ for each set B in Y .

Proof. (1) \Rightarrow (2) For $B \subseteq Y$, by the property (\mathcal{B}) , $mCl(B)$ is an m -closed set in Y . Then from Theorem 3.5 (2), it follows

$$msCl(f^{-1}(msInt(B))) \subseteq msCl(f^{-1}(msInt(mCl(B)))) \subseteq f^{-1}(mCl(B)).$$

(2) \Rightarrow (3) For $B \subseteq Y$, from Theorem 2.2 and (2),

$$\begin{aligned}
 f^{-1}(mInt(B)) &= f^{-1}(Y - mCl(Y - B)) \\
 &= X - (f^{-1}(mCl(Y - B))) \\
 &\subseteq X - msCl(f^{-1}(msInt(Y - B))) \\
 &= msInt(f^{-1}(msCl(B))).
 \end{aligned}$$

This implies $f^{-1}(mInt(B)) \subseteq msInt(f^{-1}(msCl(B)))$.

(3) \Rightarrow (1) For each m -open set V in Y , from $B = mInt(B)$ and (3), we have $f^{-1}(B) = f^{-1}(mInt(B)) \subseteq msInt(f^{-1}(msCl(B)))$. Hence f is weakly M -semicontinuous. \square

We recall the next notions: Let m_X be a minimal structure on a nonempty set X . Then (X, m_X) is said to be

- (1) m -Urysohn [6] if for each distinct points $x, y \in X$, there exist $U, V \in m_X$ containing x and y , respectively, such that $mCl(U) \cap mCl(V) = \emptyset$,
- (2) m -semi- T_2 [3] if for each distinct points $x, y \in X$, there exist m -semiopen sets U, V containing x and y , respectively, such that $U \cap V = \emptyset$.

DEFINITION 3.7. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y . Then f has a *strongly M -semiclosed graph* if for each $(x, y) \in (X \times Y) - G(f)$, there exist an m -smiopen set U containing x and an m -open set V containing y such that $(U \times mCl(V)) \cap G(f) = \emptyset$.

LEMMA 3.8. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y . Then f has a *strongly M -semiclosed graph* if and only if for each $(x, y) \in (X \times Y) - G(f)$, there exist an m -smiopen set U containing x and an m -open set V containing y such that $f(U) \cap mCl(V) = \emptyset$.

THEOREM 3.9. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y . If f is weakly M -semicontinuous and (Y, m_Y) is m -Urysohn, then $G(f)$ is a *strongly M -semiclosed graph*.

Proof. Let $(x, y) \in (X \times Y) - G(f)$; then $f(x) \neq y$. Since Y is m -Urysohn, there are m -open sets U, V containing $f(x), y$, respectively, such that $U \cap V = \emptyset$. This implies $mCl(V) \cap U = \emptyset$. On the one hand, from weak M -semicontinuity of f , for $f(x) \in U$, there exists an m -semiopen set G containing x such that $f(G) \subseteq msCl(U)$. This implies $f(G) \cap mCl(V) = \emptyset$, and by Lemma 3.8, $G(f)$ is *strongly M -semiclosed*. \square

THEOREM 3.10. Let $f : (X, m_X) \rightarrow (Y, m_Y)$ be a function on two spaces with minimal structures m_X and m_Y , respectively. If f is an *weakly M -semicontinuous injection with a strongly M -semiclosed graph*, then X is m -semi- T_2 .

Proof. Let x_1 and x_2 be any distinct points of X . Then $f(x_1) \neq f(x_2)$, so $(x_1, f(x_2)) \in (X \times Y) - G(f)$. Since the graph $G(f)$ is *strongly M -semiclosed*, there exist an m -semiopen set U containing x_1 and $V \in m_Y$ containing $f(x_2)$ such that $f(U) \cap mCl(V) = \emptyset$. Since f is weakly M -semicontinuous, for $V \in m_Y$ containing $f(x_2)$, there exists an m -semiopen set W containing x_2 in X such that $f(W) \subseteq msCl(V)$. Finally, we have $f(U) \cap f(W) = \emptyset$ because of $msCl(V) \subseteq mCl(V)$. Hence $U \cap W = \emptyset$, and X is m -semi- T_2 . \square

References

- [1] P. Monk, *An iterative finite element method for approximating the biharmonic equation*, Math. Comp. **151** (1988), 451-476.

- [2] N. Levine, *Semi-open sets and semi-continuity in topological spaces*, Ams. Math. Monthly, **70** (1963), 36-41.
- [3] W. K. Min, *m-Semiopen Sets and M-Semicontinuous Functions On Spaces With Minimal Structures*, Honam Mathematical Journal, **31**, 2009, no. 2, 239-245.
- [4] ———, *On Minimal Semicontinuous Functions*, submitted.
- [5] V. Popa and T. Noiri, *On M-continuous functions*, Anal. Univ. "Dunarea de Jos" Galati, Ser. Mat. Fiz. Mec. Teor., Fasc. II, **18** (2000), no. 23, 31-41.
- [6] ———, *On the definition of some generalized forms of continuity under minimal conditions*, Mem. Fac. Sci. Kochi. Univ. Ser. Math. **22** (2001), 9-19.
- [7] ———, *On weakly (τ, m) -continuous functions*, Rendiconti Del Circolo Matematico Di Palermo Serie II, Tomo LI, (2002), 295-316.

*

Department of Mathematics
Kangwon National University
Chuncheon 200-701, Republic of Korea
E-mail: wkmin@kangwon.ac.kr

**

Department of Mathematics
MyongJi University
Youngin 449-728, Republic of Korea.
E-mail: ykkim@mju.ac.kr