

h -STABILITY OF THE NONLINEAR PERTURBED DIFFERENTIAL SYSTEMS VIA t_∞ -SIMILARITY

YOON HOE GOO* AND SEUNG BUM YANG**

ABSTRACT. In this paper, we investigate h -stability of the nonlinear perturbed differential systems using the notion of t_∞ -similarity

1. Introduction

As is traditional in a perturbation theory of nonlinear differential equations, the behavior of solutions of a perturbed equation is determined in terms of the behavior of solutions of an unperturbed equation. There are three useful methods for studying the qualitative behavior of the solutions of perturbed nonlinear system of differential equations: the method of variation of constants formula, the second method of Lyapunov and the use of integral inequalities.

Pinto[13,15] introduced h -stability(hS) which is an important extension of the notions of exponential asymptotic stability and uniform Lipschitz stability. Also, he obtained some properties about asymptotic behavior of solutions of perturbed differential systems, some general results about asymptotic integration and gave some important examples in [14].

Choi and Ryu [3] dealt with hS of the solutions of the various differential systems and Volterra integro-differential systems. Recently, Choi et al. [4] and Goo and Ry [7,8] obtained results for hS of nonlinear differential systems via t_∞ -similarity. Goo et al.[9,10] investigated hS for the nonlinear Volterra integro-differential system.

In this paper, we investigate h -stability of the nonlinear perturbed differential systems using the notion of t_∞ -similarity.

Received May 11, 2011; Revised October 18, 2011; Accepted November 18, 2011.
2010 Mathematics Subject Classification: Primary 34D10.

Key words and phrases: h -stable, t_∞ -similarity, nonlinear nonautonomous system.
Correspondence should be addressed to Yoon Hoe Goo, yhgoo@hanseo.ac.kr.

The authors are supported by Hanseo University Fund in 2010** and 2011*.

2. Preliminaries

We consider the nonlinear nonautonomous differential system

$$(2.1) \quad x'(t) = f(t, x(t)), \quad x(t_0) = x_0,$$

where $f \in C[\mathbb{R}^+ \times \mathbb{R}^n, \mathbb{R}^n]$, $\mathbb{R}^+ = [0, \infty)$ and \mathbb{R}^n is the Euclidean n -space. We assume that the Jacobian matrix $f_x = \partial f / \partial x$ exists and is continuous on $\mathbb{R}^+ \times \mathbb{R}^n$ and $f(t, 0) = 0$.

Let $x(t) = x(t, t_0, x_0)$ denote the unique solution of (2.1) through (t_0, x_0) in $\mathbb{R}^+ \times \mathbb{R}^n$ such that $x(t_0, t_0, x_0) = x_0$. Also, we consider the associated variational systems around the zero solution of (2.1) and around $x(t)$, respectively,

$$(2.2) \quad v'(t) = f_x(t, 0)v(t), \quad v(t_0) = v_0$$

and

$$(2.3) \quad z'(t) = f_x(t, x(t, t_0, x_0))z(t), \quad z(t_0) = z_0.$$

The fundamental matrix solution $\Phi(t, t_0, x_0)$ of (2.3) is given by

$$\Phi(t, t_0, x_0) = \frac{\partial}{\partial x_0} x(t, t_0, x_0),$$

and $\Phi(t, t_0, 0)$ is the fundamental matrix solution of (2.2). The symbol $|\cdot|$ denotes arbitrary vector norm in \mathbb{R}^n .

We recall some notions of h -stability [13] and the notion of t_∞ -similarity [5].

DEFINITION 2.1. The system (2.1) (the zero solution $x = 0$ of (2.1)) is called an h -system if there exist a constant $c \geq 1$, and a positive continuous function h on \mathbb{R}^+ such that

$$|x(t)| \leq c|x_0| h(t) h(t_0)^{-1}$$

for $t \geq t_0 \geq 0$ and $|x_0|$ small enough (here $h(t)^{-1} = \frac{1}{h(t)}$).

DEFINITION 2.2. The system (2.1) (the zero solution $x = 0$ of (2.1)) is called h -stable (hS) if there exist $\delta > 0$ such that (2.1) is an h -system for $|x_0| \leq \delta$ and h is bounded.

Let \mathcal{M} denote the set of all $n \times n$ continuous matrices $A(t)$ defined on \mathbb{R}^+ and \mathcal{N} be the subset of \mathcal{M} consisting of those nonsingular matrices $S(t)$ that are of class C^1 with the property that $S(t)$ and $S^{-1}(t)$ are bounded. The notion of t_∞ -similarity in \mathcal{M} was introduced by Conti [5].

DEFINITION 2.3. A matrix $A(t) \in \mathcal{M}$ is t_∞ -similar to a matrix $B(t) \in \mathcal{M}$ if there exists an $n \times n$ matrix $F(t)$ absolutely integrable over \mathbb{R}^+ , i.e.,

$$\int_0^\infty |F(t)| dt < \infty$$

such that

$$(2.4) \quad \dot{S}(t) + S(t)B(t) - A(t)S(t) = F(t)$$

for some $S(t) \in \mathcal{N}$.

The notion of t_∞ -similarity is an equivalence relation in the set of all $n \times n$ continuous matrices on \mathbb{R}^+ , and it preserves some stability concepts [4, 11].

We give some related properties that we need in the sequel.

LEMMA 2.4. [15] *The linear system*

$$(2.5) \quad x' = A(t)x, \quad x(t_0) = x_0,$$

where $A(t)$ is an $n \times n$ continuous matrix, is an h -system (respectively h -stable) if and only if there exist $c \geq 1$ and a positive and continuous (respectively bounded) function h defined on \mathbb{R}^+ such that

$$(2.6) \quad |\phi(t, t_0)| \leq c h(t) h(t_0)^{-1}$$

for $t \geq t_0 \geq 0$, where $\phi(t, t_0)$ is a fundamental matrix of (2.5).

THEOREM 2.5. [3] *If the zero solution of (2.1) is hS, then the zero solution of (2.2) is hS.*

THEOREM 2.6. [7] *Suppose that $f_x(t, 0)$ is t_∞ -similar to $f_x(t, x(t, t_0, x_0))$ for $t \geq t_0 \geq 0$ and $|x_0| \leq \delta$ for some constant $\delta > 0$. Then the solution $v = 0$ of (2.2) is hS if and only if the solution $z = 0$ of (2.3) is hS.*

3. Main results

In this section, we investigate hS for the nonlinear perturbed differential systems.

Now, we examine the properties of hS for the perturbed system

$$(3.1) \quad y' = f(t, y) + \int_{t_0}^t g(s, y(s), Ty(s)) ds, \quad y(t_0) = y_0,$$

where $g \in C[\mathbb{R}^+ \times \mathbb{R}^n \times \mathbb{R}^n, \mathbb{R}^n]$, $g(t, 0, 0) = 0$ and $T : C(\mathbb{R}^+, \mathbb{R}^n) \rightarrow C(\mathbb{R}^+, \mathbb{R}^n)$ is a continuous operator.

THEOREM 3.1. Suppose that $f_x(t, 0)$ is t_∞ -similar to $f_x(t, x(t, t_0, x_0))$ for $t \geq t_0 \geq 0$ and $|x_0| \leq \delta$ for some constant $\delta > 0$, the solution $x = 0$ of (2.1) is hS with the increasing function h , and g in (3.1) satisfies

$$\left| \int_{t_0}^s g(\tau, y(\tau), Ty(\tau)) d\tau \right| \leq a(s)|y(s)| + b(s) \int_{t_0}^s c(\tau)|y(\tau)| d\tau, \quad t \geq t_0 \geq 0,$$

where $a, b, c \in C(\mathbb{R}^+, \mathbb{R}^+)$ and $\int_{t_0}^\infty [a(s) + b(s) \int_{t_0}^s c(\tau) d\tau] ds < \infty$. Then, the solution $y = 0$ of (3.1) is hS.

Proof. Using a slight variant of the nonlinear variation of constants formula of Alekseev[1], any solution $y(t) = y(t, t_0, x_0)$ of (3.1) passing through (t_0, x_0) is given by

$$(3.2) \quad y(t, t_0, y_0) = x(t, t_0, y_0) + \int_{t_0}^t \Phi(t, s, y(s)) \int_{t_0}^s g(\tau, y(\tau), Ty(\tau)) d\tau ds,$$

where $x(t) = x(t, t_0, x_0)$ is a solution of (2.1) passing through (t_0, x_0) . By Theorem 2.5, since the solution $x = 0$ of (2.1) is hS, the solution $v = 0$ of (2.2) is hS. Therefore, by Theorem 2.6, the solution $z = 0$ of (2.3) is hS. Applying Lemma 2.4 and the increasing property of the function h , we have

$$\begin{aligned} |y(t)| &\leq |x(t)| + \int_{t_0}^t |\Phi(t, s, y(s))| \left| \int_{t_0}^s g(\tau, y(\tau), Ty(\tau)) d\tau \right| ds \\ &\leq c_1 |y_0| h(t) h(t_0)^{-1} \\ &\quad + \int_{t_0}^t c_2 h(t) [a(s)|y(s)| h(s)^{-1} + b(s) \int_{t_0}^s c(\tau)|y(\tau)| h(\tau)^{-1} d\tau] ds. \end{aligned}$$

Set $u(t) = |y(t)| h(t)^{-1}$. Then, by Gronwall's inequality, we obtain

$$\begin{aligned} |y(t)| &\leq c_1 |y_0| h(t) h(t_0)^{-1} e^{c_2 \int_{t_0}^t [a(s) + b(s) \int_{t_0}^s c(\tau) d\tau] ds} \\ &\leq c |y_0| h(t) h(t_0)^{-1}, \quad c = c_1 e^{c_2 \int_{t_0}^\infty [a(s) + b(s) \int_{t_0}^s c(\tau) d\tau] ds} \end{aligned}$$

It follows that $y = 0$ of (3.1) is hS. Hence, the proof is complete. \square

REMARK 3.1. Letting $g(s, y(s), Ty(s)) = g(s, y(s))$, $b(s) = 0$ in Theorem 3.1, we obtain the same result as that of Theorem 3.3 in [8].

REMARK 3.2. In the linear case, we can obtain that if the zero solution $x = 0$ of (2.5) is hS, then the perturbed system

$$y' = A(t)y + \int_{t_0}^t g(s, y(s), Ty(s)) ds, \quad y(t_0) = y_0,$$

is also hS under the same hypotheses in Theorem 3.1 except the condition of t_∞ -similarity.

We need the following lemma for hS of (3.1).

LEMMA 3.2. Let $u, p, q, w, r \in C(\mathbb{R}^+, \mathbb{R}^+)$ and suppose that, for some $c \geq 0$, we have

$$(3.3) \quad u(t) \leq c + \int_{t_0}^t p(s) \int_{t_0}^s [q(\tau)u(\tau) + w(\tau) \int_{t_0}^\tau r(a)u(a)da]d\tau ds, \quad t \geq t_0.$$

Then

$$(3.4) \quad u(t) \leq c \exp\left(\int_{t_0}^t p(s) \int_{t_0}^s [q(\tau) + w(\tau) \int_{t_0}^\tau r(a)da]d\tau ds\right), \quad t \geq t_0.$$

Proof. Setting $v(t) = c + \int_{t_0}^t p(s) \int_{t_0}^s [q(\tau)u(\tau) + w(\tau) \int_{t_0}^\tau r(a)u(a)da]d\tau ds$, we have $v(t_0) = c$ and

$$(3.5) \quad \begin{aligned} v'(t) &= p(t) \int_{t_0}^t [q(s)u(s) + w(s) \int_{t_0}^s r(a)u(a)da]ds \\ &\leq p(t) \int_{t_0}^t [q(s) + w(s) \int_{t_0}^s r(a)da]v(s)ds \\ &\leq [p(t) \int_{t_0}^t [q(s) + w(s) \int_{t_0}^s r(a)da]ds]v(t), \quad t \geq t_0, \end{aligned}$$

since $v(t)$ is nondecreasing and $u(t) \leq v(t)$. It follows from the Gronwall inequality that (3.5) yields the estimate (3.4). \square

THEOREM 3.3. Suppose that $f_x(t, 0)$ is t_∞ -similar to $f_x(t, x(t, t_0, x_0))$ for $t \geq t_0 \geq 0$ and $|x_0| \leq \delta$ for some constant $\delta > 0$. If the solution $x = 0$ of (2.1) is an h -system with a positive continuous function h and g in (3.1) satisfies

$$|g(t, y, Ty)| \leq \lambda(t)|y| + \beta(t) \int_{t_0}^t \gamma(s)|y(s)|ds, \quad t \geq t_0, \quad y \in \mathbb{R}^n,$$

where $\lambda, \beta, \gamma : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ are continuous with

$$(3.6) \quad \int_{t_0}^\infty \frac{1}{h(s)} \int_{t_0}^s [h(\tau)\lambda(\tau) + \beta(\tau) \int_{t_0}^\tau h(r)\gamma(r)dr]d\tau ds < \infty,$$

for all $t_0 \geq 0$, then the solution $y = 0$ of (3.1) is an h -system.

Proof. It is known that the solutions of (2.1) and (3.1) with the same initial values are represented by the integral equation (3.2). By Theorem 2.5, since the solution $x = 0$ of (2.1) is an h -system, the solution $v = 0$

of (2.2) is an h -system. Therefore, by Theorem 2.6, the solution $z = 0$ of (2.3) is an h -system. Using (2.6) and (3.2), we obtain

$$\begin{aligned} |y(t)| &\leq |x(t)| + \int_{t_0}^t |\Phi(t, s, y(s))| \int_{t_0}^s |g(\tau, y(\tau), Ty(\tau))| d\tau ds \\ &\leq c_1 |y_0| h(t) h(t_0)^{-1} \\ &\quad + \int_{t_0}^t c_2 \frac{h(t)}{h(s)} \left[\int_{t_0}^s h(\tau) \lambda(\tau) \frac{|y(\tau)|}{h(\tau)} + \beta(\tau) \int_{t_0}^{\tau} h(r) \gamma(r) \frac{|y(r)|}{h(r)} dr d\tau \right] ds. \end{aligned}$$

Applying Lemma 3.2 with $u(t) = |y(t)| h(t)^{-1}$ and using (3.6), we obtain

$$\begin{aligned} |y(t)| &\leq c_1 |y_0| h(t) h(t_0)^{-1} e^{c_2 \int_{t_0}^t \frac{1}{h(s)} \int_{t_0}^s [h(\tau) \lambda(\tau) + \beta(\tau) \int_{t_0}^{\tau} h(r) \gamma(r) dr] d\tau ds} \\ &\leq c |y_0| h(t) h(t_0)^{-1}, \quad t \geq t_0, \end{aligned}$$

where $c = c_1 e^{c_2 \int_{t_0}^{\infty} \frac{1}{h(s)} \int_{t_0}^s [h(\tau) \lambda(\tau) + \beta(\tau) \int_{t_0}^{\tau} h(r) \gamma(r) dr] d\tau ds}$. It follows that $y = 0$ of (3.1) is an h -system. Hence, the proof is complete. \square

REMARK 3.3. In the linear case, we can obtain that if the zero solution $x = 0$ of (2.5) is an h -system, then the perturbed system

$$y' = A(t)y + \int_{t_0}^t g(s, y(s), Ty(s)) ds, \quad y(t_0) = y_0,$$

is also an h -system under the same hypotheses in Theorem 3.3 except the condition of t_∞ -similarity.

THEOREM 3.4. For the system (3.1), suppose that

$$\left| \int_{t_0}^t g(\tau, y(\tau), Ty(\tau)) d\tau \right| \leq r(t, |y|, |Ty|),$$

where $r \in C[\mathbb{R}^+ \times \mathbb{R}^+ \times \mathbb{R}^+, \mathbb{R}^+]$ is strictly increasing in u, v for each fixed $t \geq t_0 \geq 0$ with $r(t, 0, 0) = 0$. Assume also that $z = 0$ of (2.3) is hS with the nonincreasing function h . Consider the scalar differential equation

$$(3.7) \quad u' = cr(t, u, Tu), \quad u(t_0) = u_0 = c|y_0|.$$

If $u = 0$ of (3.7) is hS , then $y = 0$ of (3.1) is also hS whenever $u_0 = c|y_0|$.

Proof. Using the nonlinear variation of constants formula of Alekseev[1], the solutions of (2.1) and (3.1) with the same initial values are related by

$$y(t, t_0, y_0) = x(t, t_0, y_0) + \int_{t_0}^t \Phi(t, s, y(s)) \int_{t_0}^s g(\tau, y(\tau), Ty(\tau)) d\tau ds.$$

Then, we have

$$|y(t)| \leq |x(t)| + \int_{t_0}^t |\Phi(t, s, y(s))| \left| \int_{t_0}^s g(\tau, y(\tau), Ty(\tau)) d\tau \right| ds,$$

where $\Phi(t, s, y(s))$ is the fundamental matrix of (2.3). Then, the rest of proof can be proved in the similar manner as that of Theorem 3.4 of [8], so we omit the details. \square

Acknowledgments

The authors are very grateful for the referee's valuable comments.

References

- [1] V. M. Alekseev, *An estimate for the perturbations of the solutions of ordinary differential equations*, Vestn. Mosk. Univ. Ser. I. Math. Mekh., **2** (1961), 28–36 (Russian).
- [2] S. K. Choi and N. J. Koo, *h -stability for nonlinear perturbed systems*, Ann. of Diff. Eqs. **11** (1995), 1–9.
- [3] S. K. Choi and H. S. Ryu, *h -stability in differential systems*, Bull. Inst. Math. Acad. Sinica **21** (1993), 245–262.
- [4] S. K. Choi and N. J. Koo and H.S. Ryu, *h -stability of differential systems via t_∞ -similarity*, Bull. Korean. Math. Soc. **34** (1997), 371–383.
- [5] R. Conti, *Sulla t_∞ -similitudine tra matricie l'equivalenza asintotica dei sistemi differenziali lineari*, Rivista di Mat. Univ. Parma **8** (1957), 43–47.
- [6] S. Elaydi and R. R. M. Rao, *Lipschitz stability for nonlinear Volterra integro-differential systems*, Appl. Math. Computations **27** (1988), 191–199.
- [7] Y. H. Goo, *h -stability of the nonlinear differential systems via t_∞ -similarity*, J. Chungcheong Math. Soc. **23** (2010), 383–389.
- [8] Y. H. Goo and D. H. Ry, *h -stability of the nonlinear perturbed differential systems*, J. Chungcheong Math. Soc. **23** (2010), 827–834.
- [9] Y. H. Goo and D. H. Ry, *h -stability for perturbed integro-differential systems*, J. Chungcheong Math. Soc. **21** (2008), 511–517.
- [10] Y. H. Goo, M. H. Ji and D. H. Ry, *h -stability in certain integro-differential equations*, J. Chungcheong Math. Soc. **22** (2009), 81–88.
- [11] G. A. Hewer, *Stability properties of the equation by t_∞ -similarity*, J. Math. Anal. Appl. **41** (1973), 336–344.
- [12] V. Lakshmikantham and S. Leela, *Differential and Integral Inequalities: Theory and Applications Vol.*, Academic Press, New York and London, 1969.
- [13] M. Pinto, *Perturbations of asymptotically stable differential systems*, Analysis **4** (1984), 161–175.
- [14] M. Pinto, *Asymptotic integration of a system resulting from the perturbation of an h -system*, J. Math. Anal. Appl. **131** (1988), 194–216.
- [15] M. Pinto, *Stability of nonlinear differential systems*, Applicable Analysis **43** (1992), 1–20.

*

Department of Mathematics
Hanseo University
Seosan Chungnam 356-706, Republic of Korea
E-mail: yhgoo@hanseo.ac.kr

**

Department of Mathematics
Hanseo University
Seosan Chungnam 356-706, Republic of Korea
E-mail: ysb-good@hanmail.net