

DECOMPOSITION FORMULAS AND INTEGRAL REPRESENTATIONS FOR SOME EXTON HYPERGEOMETRIC FUNCTIONS

JUNESANG CHOI*, ANVAR HASANOV**, AND MAMASALI TURAEV***

ABSTRACT. Generalizing the Burchnell-Chaundy operator method, the authors are aiming at presenting certain decomposition formulas for the chosen six Exton functions expressed in terms of Appell's functions F_3 and F_4 , Horn's functions H_3 and H_4 , and Gauss's hypergeometric function F . We also give some integral representations for the Exton functions X_i ($i = 6, 8, 14$) each of whose kernels contains the Horn's function H_4 .

1. Introduction and preliminaries

Almost seven decades ago, Burchnell and Chaundy (see [2, 3]) and Chaundy [4] systematically presented a number of expansion and decomposition formulas for some double hypergeometric functions in series of simpler hypergeometric functions. Their method is based upon the following inverse pairs of symbolic operators:

$$\nabla_{xy}(h) := \frac{\Gamma(h)\Gamma(\delta_1 + \delta_2 + h)}{\Gamma(\delta_1 + h)\Gamma(\delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(h)_k k!}, \quad (1.1)$$

$$\Delta_{xy}(h) := \frac{\Gamma(\delta_1 + h)\Gamma(\delta_2 + h)}{\Gamma(h)\Gamma(\delta_1 + \delta_2 + h)} = \sum_{k=0}^{\infty} \frac{(-\delta_1)_k (-\delta_2)_k}{(1 - h - \delta_1 - \delta_2)_k k!}, \quad (1.2)$$

Received July 15, 2011; Accepted November 18, 2011.

2010 Mathematics Subject Classification: Primary 33C20, 33C65; Secondary 33C05, 44A45, 11Y35.

Key words and phrases: multiple hypergeometric functions, inverse pairs of symbolic operators, generalized inverse pairs of symbolic operators, Exton hypergeometric functions, Horn and Gauss hypergeometric functions, Euler type integrals, integral representations, Mellin-Barnes integrals.

Correspondence should be addressed to Junesang Choi, junesang@mail.dongguk.ac.kr.

$$\begin{aligned}
\nabla_{xy}(h) \Delta_{xy}(g) &:= \frac{\Gamma(h) \Gamma(\delta_1 + \delta_2 + h) \Gamma(\delta_1 + g) \Gamma(\delta_2 + g)}{\Gamma(\delta_1 + h) \Gamma(\delta_2 + h) \Gamma(g) \Gamma(\delta_1 + \delta_2 + g)} \\
&= \sum_{k=0}^{\infty} \frac{(g-h)_k (g)_{2k} (-\delta_1)_k (-\delta_2)_k}{(g+k-1)_k (g+\delta_1)_k (g+\delta_2)_k k!} \\
&= \sum_{k=0}^{\infty} \frac{(h-g)_k (-\delta_1)_k (-\delta_2)_k}{(h)_k (1-g-\delta_1-\delta_2)_k k!} \left(\delta_1 := x \frac{\partial}{\partial x}; \delta_2 := y \frac{\partial}{\partial y} \right).
\end{aligned} \tag{1.3}$$

In the definitions of operators (1.1) to (1.3) was used the well-known Gauss's summation theorem for the hypergeometric function ${}_2F_1 := F$ (see, e.g., [7, p. 104, Eq. (46)]):

$$\begin{aligned}
F(a, b; c; 1) &= \frac{\Gamma(c) \Gamma(c-a-b)}{\Gamma(c-a) \Gamma(c-b)} \\
&(\Re(c-a-b) > 0; c \in \mathbb{C} \setminus \mathbb{Z}_0^-),
\end{aligned} \tag{1.4}$$

where $\mathbb{C}$ is the set of complex numbers and $\mathbb{Z}_0^- := \{0, -1, -2, \dots\}$.

By employing their symbolic operators, Burchnall and Chaundy obtained several decomposition formulas for Appell's hypergeometric function (see [1, p. 14, Eqs. (11)-(14)] and, also see [7, p. 225, Eqs. (6)-(9)]) and Humbert functions of two variables, which are particularly recorded in [7, p. 94, pp. 187-188, and pp. 234-235]. The presented operators decompose the double hypergeometric functions into the product of two one-dimensional functions. Therefore, an investigation of certain properties of the double hypergeometric functions can be made by means of one-dimensional functions. It is interesting to note that this method has been forgotten without any particular reason, though this method turned out to be useful in application of the other hypergeometric functions. We note that in monograph [25] was given the definitions of 205 hypergeometric functions with their regions of convergence. Furthermore, in the most recent works [5, 11, 12, 13, 14], aforementioned two-dimensional operators (1.1) to (1.3) were generalized to the following multi-dimensional inverse pairs of symbolic operators:

$$\begin{aligned}
\tilde{\nabla}_{x_1; x_2, \dots, x_r}(h) &:= \frac{\Gamma(h) \Gamma(\delta_1 + \delta_2 + \dots + \delta_r + h)}{\Gamma(\delta_1 + h) \Gamma(\delta_2 + \dots + \delta_r + h)} \\
&= \sum_{m_2, \dots, m_r=0}^{\infty} \frac{(-\delta_1)_{m_2+\dots+m_r} (-\delta_2)_{m_2} \dots (-\delta_r)_{m_r}}{(h)_{m_2+\dots+m_r} m_2! \dots m_r!},
\end{aligned} \tag{1.5}$$

$$\begin{aligned}
\tilde{\Delta}_{x_1; x_2, \dots, x_r}(h) &:= \frac{\Gamma(\delta_1 + h) \Gamma(\delta_2 + \dots + \delta_r + h)}{\Gamma(h) \Gamma(\delta_1 + \delta_2 + \dots + \delta_r + h)} \\
&= \sum_{m_2, \dots, m_r=0}^{\infty} \frac{(-\delta_1)_{m_2+\dots+m_r} (-\delta_2)_{m_2} \dots (-\delta_r)_{m_r}}{(1-h-\delta_1-\dots-\delta_r)_{m_2+\dots+m_r} m_2! \dots m_r!} \\
&= \sum_{m_2, \dots, m_r=0}^{\infty} \frac{(-1)^{m_2+\dots+m_r}}{(h+m_2+\dots+m_r-1)_{m_2+\dots+m_r} (h+\delta_1)_{m_2+\dots+m_r}} \\
&\quad \cdot \frac{(h)_{2(m_2+\dots+m_r)} (-\delta_1)_{m_2+\dots+m_r} (-\delta_2)_{m_2} \dots (-\delta_r)_{m_r}}{(h+\delta_2+\dots+\delta_r)_{m_2+\dots+m_r} m_2! \dots m_r!}
\end{aligned} \tag{1.6}$$

$$\left(\delta_j := x_j \frac{\partial}{\partial x_j}; j = 1, 2, \dots, r \right).$$

By means of generalized symbolic operators (1.5) and (1.6), decomposition formulas for Lauricella function (see [1, p. 114, Eqs. (1)-(4)]), Horn functions (see [1, p. 224, Eqs. (10)-(11)], and Srivastava's hypergeometric functions (see [23, 24], also see [25, p. 43, Eqs. (11)-(13)]) have been presented. It should be noted that the obtained decomposition formulas have then been used in the solution of boundary-value problems of degenerated differential equations of second order (see [9, 10, 15, 22]). An attentive reader may have observed that there is no inverse pairs of symbolic operators that can be used in one-dimensional case. For this purpose we define following operators:

$$\begin{aligned}
H_{x_1, \dots, x_l}(\alpha, \beta) &:= \frac{\Gamma(\beta) \Gamma(\alpha + \delta_1 + \dots + \delta_l)}{\Gamma(\alpha) \Gamma(\beta + \delta_1 + \dots + \delta_l)} \\
&= \sum_{k_1, \dots, k_l=0}^{\infty} \frac{(\beta - \alpha)_{k_1+\dots+k_l} (-\delta_1)_{k_1} \dots (-\delta_l)_{k_l}}{(\beta)_{k_1+\dots+k_l} k_1! \dots k_l!}
\end{aligned} \tag{1.7}$$

and

$$\begin{aligned}
\bar{H}_{x_1, \dots, x_l}(\alpha, \beta) &:= \frac{\Gamma(\alpha) \Gamma(\beta + \delta_1 + \dots + \delta_l)}{\Gamma(\beta) \Gamma(\alpha + \delta_1 + \dots + \delta_l)} \\
&= \sum_{k_1, \dots, k_l=0}^{\infty} \frac{(\beta - \alpha)_{k_1+\dots+k_l} (-\delta_1)_{k_1} \dots (-\delta_l)_{k_l}}{(1 - \alpha - \delta_1 - \dots - \delta_l)_{k_1+\dots+k_l} k_1! \dots k_l!} \\
&\quad \left(\delta_j := x_j \frac{\partial}{\partial x_j}, j = 1, \dots, l; l \in \mathbb{N} := \{1, 2, 3, \dots\} \right).
\end{aligned} \tag{1.8}$$

In order to obtain aforementioned operators, we have applied some known multiple hypergeometric summation formulas such as (see [2,

p. 117]):

$$F_D^{(r)}(\alpha; \beta_1, \dots, \beta_r; \gamma; 1, \dots, 1) = \frac{\Gamma(\gamma) \Gamma(\gamma - \alpha - \beta_1 - \dots - \beta_r)}{\Gamma(\gamma - \alpha) \Gamma(\gamma - \beta_1 - \dots - \beta_r)}$$

$$(\Re(\gamma - \alpha - \beta_1 - \dots - \beta_r) > 0; \gamma \in \mathbb{C} \setminus \mathbb{Z}_0^-)$$

for Lauricella function $F_D^{(r)}$ in r variables (see [1, p. 117]).

Here, applying the inverse pairs of operators (1.7) and (1.8) to Exton functions X_i ($i = 2, 6, 8, 14, 17, 20$) recalled below among his twenty ones (see [8], also see [25, pp. 113–114]), we aim at presenting certain decomposition formulas for the chosen Exton functions expressed in terms of Appell's functions F_3 and F_4 , Horn's functions H_3 and H_4 , and Gauss's hypergeometric function F . We also give some integral representations for the Exton functions X_i ($i = 6, 8, 14$) each of whose kernels contains the Horn's function H_4 .

We recall, for convenience, the six Exton functions:

$$X_2(a_1, a_2; c_1, c_2, c_3; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+2n+p} (a_2)_p}{(c_1)_m (c_2)_n (c_3)_p m!n!p!} x^m y^n z^p$$

$$(\{2\sqrt{r} + 2\sqrt{s} + 1 < 1\}, |x| \leq r, |y| \leq s, |z| \leq t); \quad (1.9)$$

$$X_6(a_1, a_2, a_3; c_1, c_2; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n+p} (a_2)_n (a_3)_p}{(c_1)_{m+n} (c_2)_p m!n!p!} x^m y^n z^p$$

$$(\{t + 2\sqrt{r} < 1 \wedge s < \frac{1}{2}(1-t) + \frac{1}{2}\sqrt{(1-t)^2 - 4r}\},$$

$$|x| \leq r, |y| \leq s, |z| \leq t); \quad (1.10)$$

$$X_8(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n+p} (a_2)_n (a_3)_p}{(c_1)_m (c_2)_n (c_3)_p m!n!p!} x^m y^n z^p$$

$$(\{2\sqrt{r} + s + t < 1\}, |x| \leq r, |y| \leq s, |z| \leq t); \quad (1.11)$$

$$X_{14}(a_1, a_2, a_3; c_1, c_2; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{n+p} (a_3)_p}{(c_1)_{m+n} (c_2)_p m!n!p!} x^m y^n z^p$$

$$(1.12)$$

$$\left(\left\{ r < \frac{1}{4} \wedge t < 1 \wedge s < (1-t) \left[\frac{1}{2} + \frac{1}{2} \sqrt{1-4r} \right] \right\}, \right. \\ \left. |x| \leq r, \quad |y| \leq s, \quad |z| \leq t \right);$$

$$X_{15}(a_1, a_2, a_3; c_1, c_2; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{n+p} (a_3)_p}{(c_1)_m (c_2)_{n+p} m!n!p!} x^m y^n z^p \\ (\{s + 2\sqrt{r} < 1 \wedge t < 1\}, \quad |x| \leq r, \quad |y| \leq s, \quad |z| \leq t); \quad (1.13)$$

$$X_{17}(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_{n+p} (a_3)_p}{(c_1)_m (c_2)_n (c_3)_p m!n!p!} x^m y^n z^p \\ (1.14)$$

$$\left(\left\{ r < \frac{1}{4} \wedge t < 1 \wedge s < (1-2\sqrt{r})(1-t) \right\}, \quad |x| \leq r, \quad |y| \leq s, \quad |z| \leq t \right);$$

$$X_{20}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z) \\ = \sum_{m,n,p=0}^{\infty} \frac{(a_1)_{2m+n} (a_2)_n (a_3)_p (a_4)_p}{(c_1)_{m+p} (c_2)_n m!n!p!} x^m y^n z^p \\ (\{s + 2\sqrt{r} < 1 \wedge t < 1\}, \quad |x| \leq r, \quad |y| \leq s, \quad |z| \leq t). \quad (1.15)$$

It is *remarked* in passing that the Exton functions have recently been actively investigated (see, e.g., [6, 16, 17, 18, 19]).

2. A set of operator identities

By using Burchnall and Chaundy [2, 3] and Chaundy [4] method together with symbolic operators (1.7) and (1.8), we find the following set of operator identities for hypergeometric Exton functions:

$$X_2(a_1, a_2; c_1, c_2, c_3; x, y, z) = H_z(a_2, c_3) \\ \cdot (1-z)^{-a_1} F_4 \left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1, c_2; \frac{4x}{(1-z)^2}, \frac{4y}{(1-z)^2} \right); \quad (2.1)$$

$$(1-z)^{-a_1} F_4 \left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1, c_2; \frac{4x}{(1-z)^2}, \frac{4y}{(1-z)^2} \right) \\ = \bar{H}_z(a_2, c_3) X_2(a_1, a_2; c_1, c_2, c_3; x, y, z); \quad (2.2)$$

$$\begin{aligned} X_6(a_1, a_2, a_3; c_1, c_2; x, y, z) \\ = H_z(a_3, c_2)(1-z)^{-a_1} H_3\left(a_1, a_2; c_1; \frac{x}{(1-z)^2}, \frac{y}{1-z}\right); \end{aligned} \quad (2.3)$$

$$\begin{aligned} (1-z)^{-a_1} H_3\left(a_1, a_2; c_1; \frac{x}{(1-z)^2}, \frac{y}{1-z}\right) \\ = \bar{H}_z(a_3, c_2) X_6(a_1, a_2, a_3; c_1, c_2; x, y, z); \end{aligned} \quad (2.4)$$

$$\begin{aligned} X_8(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) \\ = H_z(a_3, c_3)(1-z)^{-a_1} H_4\left(a_1, a_2; c_1, c_2; \frac{x}{(1-z)^2}, \frac{y}{1-z}\right); \end{aligned} \quad (2.5)$$

$$\begin{aligned} (1-z)^{-a_1} H_4\left(a_1, a_2; c_1, c_2; \frac{x}{(1-z)^2}, \frac{y}{1-z}\right) \\ = \bar{H}_z(a_3, c_3) X_8(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z); \end{aligned} \quad (2.6)$$

$$\begin{aligned} X_{14}(a_1, a_2, a_3; c_1, c_2; x, y, z) \\ = H_z(a_3, c_2)(1-z)^{-a_2} H_3\left(a_1, a_2; c_1; x, \frac{y}{1-z}\right); \end{aligned} \quad (2.7)$$

$$\begin{aligned} (1-z)^{-a_2} H_3\left(a_1, a_2; c_1; x, \frac{y}{1-z}\right) \\ = \bar{H}_z(a_3, c_2) X_{14}(a_1, a_2, a_3; c_1, c_2; x, y, z); \end{aligned} \quad (2.8)$$

$$\begin{aligned} X_{15}(a_1, a_2, a_3; c_1, c_2; x, y, z) \\ = H_{y,z}(a_2, c_2)(1-y)^{-a_1}(1-z)^{-a_3} F\left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1; \frac{4x}{(1-y)^2}\right); \end{aligned} \quad (2.9)$$

$$\begin{aligned} (1-y)^{-a_1}(1-z)^{-a_3} F\left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1; \frac{4x}{(1-y)^2}\right) \\ = \bar{H}_{y,z}(a_2, c_2) X_{15}(a_1, a_2, a_3; c_1, c_2; x, y, z); \end{aligned} \quad (2.10)$$

$$\begin{aligned} X_{17}(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) \\ = H_z(a_3, c_3)(1-z)^{-a_2} H_4\left(a_1, a_2; c_1, c_2; x, \frac{y}{1-z}\right); \end{aligned} \quad (2.11)$$

$$\begin{aligned} (1-z)^{-a_2} H_4\left(a_1, a_2; c_1, c_2; x, \frac{y}{1-z}\right) \\ = \bar{H}_z(a_3, c_3) X_{17}(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z); \end{aligned} \quad (2.12)$$

$$\begin{aligned}
& X_{20}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z) \\
&= H_y(a_2, c_2)(1-y)^{-a_1} F_3\left(\frac{a_1}{2}, a_3, \frac{a_1}{2} + \frac{1}{2}, a_4; c_1; \frac{4x}{(1-y)^2}, z\right); \tag{2.13}
\end{aligned}$$

$$\begin{aligned}
& (1-y)^{-a_1} F_3\left(\frac{a_1}{2}, a_3, \frac{a_1}{2} + \frac{1}{2}, a_4; c_1; \frac{4x}{(1-y)^2}, z\right) \\
&= \bar{H}_y(a_2, c_2) X_{20}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z); \tag{2.14}
\end{aligned}$$

where

$$\begin{aligned}
& F_3(a_1, a_2, b_1, b_2; c; x, y) \\
&= \sum_{m,n=0}^{\infty} \frac{(a_1)_m (a_2)_n (b_1)_m (b_2)_n}{(c)_{m+n} m! n!} x^m y^n \quad (|x| < 1, |y| < 1)
\end{aligned}$$

and

$$F_4(a, b; c_1, c_2; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{m+n} (b)_{m+n}}{(c_1)_m (c_2)_n m! n!} x^m y^n \quad (\sqrt{|x|} + \sqrt{|y|} < 1)$$

are Appell's hypergeometric functions (see [7, p. 224, Eqs. (8) and (9)]) and

$$H_3(a, b; c; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m+n} (b)_n}{(c)_{m+n} m! n!} x^m y^n \quad \left(|x| + \left(|y| - \frac{1}{2}\right)^2 \leq \frac{1}{4}\right)$$

and

$$H_4(a, b; c_1, c_2; x, y) = \sum_{m,n=0}^{\infty} \frac{(a)_{2m+n} (b)_n}{(c_1)_m (c_2)_n m! n!} x^m y^n \quad (4|x| \leq (|y| - 1)^2)$$

are the Horn's hypergeometric functions (see [7, p. 225, Eqs. (15) and (16)]), and F is Gauss hypergeometric function (see [7, p. 56, Eq. (1)]). The proof of the operator identities (2.1) to (2.14) is based on employing of Mellin [20], and Mellin-Barnes integral representation methods (see [7, p. 232]) for hypergeometric functions. The details involved in these derivations are omitted here.

3. Decomposition formulas

In [21, p. 93], it is proved that, for any analytic functions f , the following equalities hold true:

$$(\delta_\xi + \alpha)_n \{f(\xi)\} = \xi^{1-\alpha} \frac{d^n}{d\xi^n} \{\xi^{\alpha+n-1} f(\xi)\} \tag{3.1}$$

$$\left(\delta_\xi := \xi \frac{d}{d\xi}; \alpha \in \mathbb{C}; n \in \mathbb{N}_0 := \mathbb{N} \cup \{0\} \right)$$

and

$$(-\delta_\xi)_n \{f(\xi)\} = (-1)^n \xi^n \frac{d^n}{d\xi^n} \{f(\xi)\} \quad \left(\delta_\xi := \xi \frac{d}{d\xi}; n \in \mathbb{N}_0 \right). \quad (3.2)$$

In view of formulas (3.1) and (3.2), and taking into account the differentiation formula for hypergeometric functions, from operator identities (2.1) to (2.14), we find the following decomposition formulas:

$$\begin{aligned} X_2(a_1, a_2; c_1, c_2, c_3; x, y, z) &= (1-z)^{-a_1} \sum_{i=0}^{\infty} \frac{(a_1)_i (c_3 - a_2)_i}{(c_3)_i i!} \left(\frac{z}{z-1} \right)^i \\ &\quad \cdot F_4 \left(\frac{a_1+i}{2}, \frac{a_1+i}{2} + \frac{1}{2}; c_1, c_2; \frac{4x}{(1-z)^2}, \frac{4y}{(1-z)^2} \right); \end{aligned} \quad (3.3)$$

$$\begin{aligned} &(1-z)^{-a_1} F_4 \left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1, c_2; \frac{4x}{(1-z)^2}, \frac{4y}{(1-z)^2} \right) \\ &= \sum_{i=0}^{\infty} \frac{(a_1)_i (c_3 - a_2)_i}{(c_3)_i i!} z^i X_2(a_1+i, a_2; c_1, c_2, c_3+i; x, y, z); \end{aligned} \quad (3.4)$$

$$\begin{aligned} X_6(a_1, a_2, a_3; c_1, c_2; x, y, z) &= (1-z)^{-a_1} \sum_{i=0}^{\infty} \frac{(-1)^i (a_1)_i (c_2 - a_3)_i}{(c_2)_i i!} \\ &\quad \cdot \left(\frac{z}{1-z} \right)^i H_3 \left(a_1+i, a_2; c_1; \frac{x}{(1-z)^2}, \frac{y}{1-z} \right); \end{aligned} \quad (3.5)$$

$$\begin{aligned} &(1-z)^{-a_1} H_3 \left(a_1, a_2; c_1; \frac{x}{(1-z)^2}, \frac{y}{1-z} \right) \\ &= \sum_{i=0}^{\infty} \frac{(a_1)_i (c_2 - a_3)_i (-p)_i}{(c_2)_i i!} z^i X_6(a_1+i, a_2, a_3; c_1, c_2+i; x, y, z); \end{aligned} \quad (3.6)$$

$$\begin{aligned} X_8(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) &= (1-z)^{-a_1} \sum_{i=0}^{\infty} \frac{(-1)^i (a_1)_i (c_3 - a_3)_i}{(c_3)_i i!} \\ &\quad \cdot \left(\frac{z}{1-z} \right)^i H_4 \left(a_1+i, a_2; c_1, c_2; \frac{x}{(1-z)^2}, \frac{y}{1-z} \right); \end{aligned} \quad (3.7)$$

$$\begin{aligned}
& (1-z)^{-a_1} H_4 \left(a_1, a_2; c_1, c_2; \frac{x}{(1-z)^2}, \frac{y}{1-z} \right) \\
&= \sum_{i=0}^{\infty} \frac{(a_1)_i (c_3 - a_3)_i}{(c_3)_i i!} z^i X_8(a_1 + i, a_2, a_3; c_1, c_2, c_3 + i; x, y, z);
\end{aligned} \tag{3.8}$$

$$\begin{aligned}
X_{14}(a_1, a_2, a_3; c_1, c_2; x, y, z) &= (1-z)^{-a_2} \sum_{i=0}^{\infty} \frac{(-1)^i (a_2)_i (c_2 - a_3)_i}{(c_2)_i i!} \\
&\quad \cdot \left(\frac{z}{1-z} \right)^i H_3 \left(a_1, a_2 + i; c_1; x, \frac{y}{1-z} \right);
\end{aligned} \tag{3.9}$$

$$\begin{aligned}
& (1-z)^{-a_2} H_3 \left(a_1, a_2; c_1; x, \frac{y}{1-z} \right) \\
&= \sum_{i=0}^{\infty} \frac{(a_2)_i (c_2 - a_3)_i}{(c_2)_i i!} z^i X_{14}(a_1, a_2 + i, a_3; c_1, c_2 + i; x, y, z);
\end{aligned} \tag{3.10}$$

$$\begin{aligned}
& X_{15}(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= (1-y)^{-a_1} (1-z)^{-a_3} \sum_{i,j=0}^{\infty} \frac{(-1)^{i+j} (a_1)_i (a_3)_j (c_2 - a_2)_{i+j}}{(c_2)_{i+j} i! j!} \\
&\quad \cdot \left(\frac{y}{1-y} \right)^i \left(\frac{z}{1-z} \right)^j F \left(\frac{a_1 + i}{2}, \frac{a_1 + i}{2} + \frac{1}{2}; c_1; \frac{4x}{(1-y)^2} \right);
\end{aligned} \tag{3.11}$$

$$\begin{aligned}
& (1-y)^{-a_1} (1-z)^{-a_3} F \left(\frac{a_1}{2}, \frac{a_1}{2} + \frac{1}{2}; c_1; \frac{4x}{(1-y)^2} \right) \\
&= \sum_{i,j=0}^{\infty} \frac{(a_1)_i (a_3)_j (c_2 - a_2)_{i+j}}{(c_2)_{i+j} i! j!} y^i z^j \\
&\quad \cdot X_{15}(a_1 + i, a_2, a_3 + j; c_1, c_2 + i + j; x, y, z);
\end{aligned} \tag{3.12}$$

$$\begin{aligned}
& X_{17}(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) \\
&= (1-z)^{-a_2} \sum_{i=0}^{\infty} \frac{(-1)^i (a_2)_i (c_3 - a_3)_i}{(c_3)_i i!} \left(\frac{z}{1-z} \right)^i \\
&\quad \cdot H_4 \left(a_1, a_2 + i; c_1, c_2; x, \frac{y}{1-z} \right);
\end{aligned} \tag{3.13}$$

$$\begin{aligned}
& (1-z)^{-a_2} H_4 \left(a_1, a_2; c_1, c_2; x, \frac{y}{1-z} \right) \\
&= \sum_{i=0}^{\infty} \frac{(a_2)_i (c_3 - a_3)_i}{(c_3)_i i!} z^i X_{17}(a_1, a_2 + i, a_3; c_1, c_2, c_3 + i; x, y, z);
\end{aligned} \tag{3.14}$$

$$\begin{aligned}
X_{20}(a_1, a_2, a_3, a_4; c_1, c_2; x, y, z) &= (1-y)^{-a_1} \sum_{i=0}^{\infty} \frac{(-1)^i (a_1)_i (c_2 - a_2)_i}{(c_2)_i i!} \\
&\quad \cdot \left(\frac{y}{1-y} \right)^i F_3 \left(\frac{a_1 + i}{2}, a_3, \frac{a_1 + i}{2} + \frac{1}{2}, a_4; c_1; \frac{4x}{(1-y)^2}, z \right);
\end{aligned} \tag{3.15}$$

$$\begin{aligned}
& (1-y)^{-a_1} F_3 \left(\frac{a_1}{2}, a_3, \frac{a_1}{2} + \frac{1}{2}, a_4; c_1; \frac{4x}{(1-y)^2}, z \right) \\
&= \sum_{i=0}^{\infty} \frac{(a_1)_i (c_2 - a_2)_i}{(c_2)_i i!} y^i X_{20}(a_1 + i, a_2, a_3, a_4; c_1, c_2 + i; x, y, z).
\end{aligned} \tag{3.16}$$

4. Integral representations

We present certain interesting integral representations.

$$\begin{aligned}
H_3(a, b; c; x, y) &= \frac{\Gamma(c)}{\Gamma(b)\Gamma(c-b)} \int_0^1 \xi^{b-1} (1-\xi)^{c-b-1} (1-y\xi)^{-a} \\
&\quad \cdot F \left(\frac{a}{2}, \frac{1}{2} + \frac{a}{2}; c-b; \frac{4x(1-\xi)}{(1-y\xi)^2} \right) d\xi \quad (\Re(c) > \Re(b) > 0);
\end{aligned} \tag{4.1}$$

$$\begin{aligned}
& H_4(a, b; c_1, c_2; x, y) \\
&= \frac{\Gamma(c_2)}{\Gamma(b)\Gamma(c_2-b)} \int_0^1 \xi^{b-1} (1-\xi)^{c_2-b-1} (1-y\xi)^{-a} \\
&\quad \cdot F\left(\frac{a}{2}, \frac{1}{2} + \frac{a}{2}; c_1; \frac{4x}{(1-y\xi)^2}\right) d\xi \quad (\Re(c_2) > \Re(b) > 0);
\end{aligned} \tag{4.2}$$

$$\begin{aligned}
& X_6(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= \frac{\Gamma(c_1)}{\Gamma(a_2)\Gamma(c_1-a_2)} \int_0^1 \xi^{a_2-1} (1-\xi)^{c_1-a_2-1} (1-z-y\xi)^{-a_1} \\
&\quad \cdot H_4\left(a_1, c_2-a_3; c_1-a_2, c_2; \frac{x(1-\xi)}{(1-z-y\xi)^2}, -\frac{z}{1-z-y\xi}\right) d\xi \\
&\quad (\Re(c_1) > \Re(a_2) > 0);
\end{aligned} \tag{4.3}$$

$$\begin{aligned}
& X_8(a_1, a_2, a_3; c_1, c_2, c_3; x, y, z) \\
&= \frac{\Gamma(c_2)}{\Gamma(a_2)\Gamma(c_2-a_2)} \int_0^1 \xi^{a_2-1} (1-\xi)^{c_2-a_2-1} (1-z-y\xi)^{-a_1} \\
&\quad \cdot H_4\left(a_1, c_3-a_3; c_1, c_3; \frac{x}{(1-z-y\xi)^2}, -\frac{z}{1-z-y\xi}\right) d\xi \\
&\quad (\Re(c_2) > \Re(a_2) > 0);
\end{aligned} \tag{4.4}$$

$$\begin{aligned}
& X_{14}(a_1, a_2, a_3; c_1, c_2; x, y, z) \\
&= (1-z)^{-a_2} \frac{\Gamma(a_1+a_2)}{\Gamma(a_1)\Gamma(a_2)} \int_0^1 \xi^{a_1-1} (1-\xi)^{a_2-1} \\
&\quad \cdot H_4\left(a_1+a_2, c_2-a_3; c_1, c_2; x\xi^2 + \frac{y\xi(1-\xi)}{1-z}, -\frac{z(1-\xi)}{1-z}\right) d\xi \\
&\quad (\Re(a_1) > 0; \Re(a_2) > 0).
\end{aligned} \tag{4.5}$$

Proof. It is noted that each of the integral representations in Section 4 can be proved mainly by expressing the series definition of the involved special function in each integrand and changing the order of the integral sign and the summation, and finally using the following well-known relationship between the Beta function $B(\alpha, \beta)$ and the Gamma function

Γ :

$$(4.6) \quad B(\alpha, \beta) := \begin{cases} \int_0^1 t^{\alpha-1}(1-t)^{\beta-1} dt & (\Re(\alpha) > 0; \Re(\beta) > 0) \\ \frac{\Gamma(\alpha)\Gamma(\beta)}{\Gamma(\alpha+\beta)} & (\alpha, \beta \in \mathbb{C} \setminus \mathbb{Z}_0^-). \end{cases}$$

□

References

- [1] P. Appell and J. Kampé de Fériet, *Fonctions Hypergeometriques et Hyperspheriques; Polynomes d'Hermite*, Gauthier - Villars, Paris, 1926.
- [2] J. L. Burchnall and T. W. Chaundy, Expansions of Appell's double hypergeometric functions, *Quart. J. Math. Oxford Ser.* **11** (1940), 249–270.
- [3] J. L. Burchnall and T. W. Chaundy, Expansions of Appell's double hypergeometric functions. II, *Quart. J. Math. Oxford Ser.* **12** (1941), 112–128.
- [4] T. W. Chaundy, Expansions of hypergeometric functions, *Quart. J. Math. Oxford Ser.* **13** (1942), 159–171.
- [5] J. Choi and A. Hasanov, Applications of the operator $H(\alpha, \beta)$ to the Humbert double hypergeometric functions, *Comput. Math. Appl.* **61** (2011), 663–671.
- [6] J. Choi, A. K. Rathie and H. Harsh, Remarks on a summation formula for three variables hypergeometric function and certain hypergeometric transformations, *East Asian Math. J.* **25** (4) (2009), 481–486.
- [7] A. Erdélyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions*, Vol. **I**, McGraw-Hill Book Company, New York, Toronto and London, 1953.
- [8] H. Exton, Hypergeometric functions of three variables, *J. Indian Acad. Math.* **4** (1982), 113–119.
- [9] A. Hasanov, Fundamental solutions of generalized bi-axially symmetric Helmholtz equation, *Complex Variables and Elliptic Equations* **52** (2007), no. 8, 673–683.
- [10] A. Hasanov and E. T. Karimov, Fundamental solutions for a class of three-dimensional elliptic equations with singular coefficients, *Appl. Math. Lett.* **22** (2009), 1828–1832.
- [11] A. Hasanov and H. M. Srivastava, Decomposition formulas associated with the Lauricella multivariable hypergeometric functions, *Comput. Math. Appl.* **53** (2007), 1119–1128.
- [12] A. Hasanov and H. M. Srivastava, Some decomposition formulas associated with the Lauricella function $F_A^{(r)}$ and other multiple hypergeometric functions, *Appl. Math. Lett.* **19** (2006), 113–121.
- [13] A. Hasanov, H. M. Srivastava, and M. Turaev, Decomposition formulas for some triple hypergeometric functions, *J. Math. Anal. Appl.* **324** (2006), 955–969.
- [14] A. Hasanov and M. Turaev, Decomposition formulas for the double hypergeometric functions G_1 and G_2 , *Appl. Math. Comput.* **187** (2007), 195–201.

- [15] E. T. Karimov, On a boundary problem with Neumann's condition for 3D singular elliptic equations, *Appl. Math. Lett.* **23** (2010), 517–522.
- [16] Y. S. Kim, J. Choi and A. K. Rathie, Remark on two results by Padmanabham for Exton's triple hypergeometric series, *Honam Math. J.* **27** (2005), no. 4, 603–608.
- [17] Y. S. Kim and A. K. Rathie, On extension formulas for the triple hypergeometric series due to Exton, *Bull. Korean Math. Soc.* **44** (2007), no. 4, 743–751.
- [18] Y. S. Kim and A. K. Rathie, Another method for Padmanabham's transformation formula for Exton's triple hypergeometric series, *Commun. Korean Math. Soc.* **24** (2009), no. 4, 517–521.
- [19] S. W. Lee and Y. S. Kim, An extension of the triple hypergeometric series by Exton, *Honam Math. J.* **32** (2010), no. 1, 61–71.
- [20] O. I. Marichev, *Handbook of Integral Transforms of Higher Transcendental Functions: Theory and algorithmic Tables*, Halsted Press (Ellis Horwood Limited, Chichester), Wiley, New York, Chichester, Brisbane and Toronto, 1982.
- [21] E. G. Poole, *Introduction to the Theory of Linear Differential Equations*, Clarendon (Oxford University Press), Oxford, 1936.
- [22] M. S. Salakhitdinov and A. Hasanov, A solution of the Neumann-Dirichlet boundary value problem for generalized bi-axially symmetric Helmholtz equation, *Complex Variables and Elliptic Equations* **53** (2008), no. 4, 355–364.
- [23] H. M. Srivastava, Hypergeometric functions of three variables, *Ganita* **15** (1964), 97–108.
- [24] H. M. Srivastava, Some integrals representing triple hypergeometric functions, *Rend. Circ. Mat. Palermo Ser. 2* **16** (1967), 99–115.
- [25] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Halsted Press (Ellis Horwood Limited, Chichester), Wiley, New York, Chichester, Brisbane, and Toronto, 1985.

*

Department of Mathematics
Dongguk University
Gyeongju 780-714, Republic of Korea
E-mail: junesang@mail.dongguk.ac.kr

**

Department of Mathematics
Dongguk University
Gyeongju 780-714, Republic of Korea
E-mail: ahasanov@dongguk.ac.kr

Department of Mathematics
Dongguk University
Gyeongju 780-714, Republic of Korea
E-mail: mturaev@dongguk.ac.kr